



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification n <sup>5</sup> : C12Q 1/68, C12P 19/34 C07H 15/12, G01N 33/53	A1	(11) International Publication Number: WO 91/01384 (43) International Publication Date: 7 February 1991 (07.02.91)
--	----	---

(21) International Application Number: PCT/US90/03907

(22) International Filing Date: 10 July 1990 (10.07.90)

(30) Priority data:  
379,501 11 July 1989 (11.07.89) US(71) Applicant: GEN-PROBE, INCORPORATED [US/US];  
9880 Campus Point Drive, San Diego, CA 92121 (US).(72) Inventors: KACIAN, Daniel, Louis ; 3911 Tambor Road,  
San Diego, CA 92124 (US). FULTZ, Timothy, J. ; 1872  
Teakwood Way, Vista, CA 92083 (US).(74) Agents: CONSALVI, Mary, S. et al.; 611 West Sixth Street,  
34th Floor, Los Angeles, CA 90017 (US).


(81) Designated States: AU, FI, JP, KR, NO.

**Published***With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: NUCLEIC ACID SEQUENCE AMPLIFICATION METHODS UTILIZING A TRANSCRIPTION COMPLEX

## (57) Abstract

Methods of synthesizing multiple copies of a target nucleic acid sequence autocatalytically under conditions of substantially constant temperature, ionic strength, and pH are provided in which multiple RNA copies of the target sequence autocatalytically generate additional copies. These methods are useful for generating copies of a nucleic acid target sequence for purposes which include assays to quantitate specific nucleic acid sequences in clinical, environmental, forensic and similar samples, cloning and generating probes.



### DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

#### *FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
DE	Germany	LU	Luxembourg	TD	Chad
DK	Denmark			TG	Togo
				US	United States of America

DESCRIPTION

Nucleic Acid Sequence Amplification Methods Utilizing a Transcription Complex.

This application is a continuation-in-part of App. Ser. No. 379,501 filed July 11, 1989.

Field of the Invention

5        This invention relates to methods for increasing the number of copies of a specific nucleic acid sequence or "target sequence" which may be present either alone or as a component, large or small, of a homogeneous or heterogeneous mixture of nucleic acids. The mixture of nucleic  
10 acids may be that found in a sample taken for diagnostic testing, environmental testing, for research studies, for the preparation of reagents or materials for other processes such as cloning, or for other purposes.

      The selective amplification of specific nucleic acid  
15 sequences is of value in increasing the sensitivity of diagnostic and environmental assays while maintaining specificity; increasing the sensitivity, convenience, accuracy and reliability of a variety of research procedures; and providing ample supplies of specific  
20 oligonucleotides for various purposes.

      The present invention is particularly suitable for use in environmental and diagnostic testing due to the convenience with which it may be practiced.

25 Background of the Invention

      The detection and/or quantitation of specific nucleic acid sequences is an increasingly important technique for identifying and classifying microorganisms, diagnosing infectious diseases, detecting and characterizing genetic  
30 abnormalities, identifying genetic changes associated with cancer, studying genetic susceptibility to disease, and measuring response to various types of treatment. Such procedures have also found expanding uses in detecting and

quantitating microorganisms in foodstuffs, environmental samples, seed stocks, and other types of material where the presence of specific microorganisms may need to be monitored. Other applications are found in the forensic sciences, anthropology, archaeology, and biology where measurement of the relatedness of nucleic acid sequences has been used to identify criminal suspects, resolve paternity disputes, construct genealogical and phylogenetic trees, and aid in classifying a variety of life forms.

A common method for detecting and quantitating specific nucleic acid sequences is nucleic acid hybridization. This method is based on the ability of two nucleic acid strands which contain complementary or essentially complementary sequences to specifically associate, under appropriate conditions, to form a double-stranded structure. To detect and/or quantitate a specific nucleic acid sequence (known as the "target sequence"), a labelled oligonucleotide (known as a "probe") is prepared which contains sequences complementary to those of the target sequence. The probe is mixed with a sample suspected of containing the target sequence, and conditions suitable for hybrid formation are created. The probe hybridizes to the target sequence if it is present in the sample. The probe-target hybrids are then separated from the single-stranded probe in one of a variety of ways. The amount of label associated with the hybrids is measured.

The sensitivity of nucleic acid hybridization assays is limited primarily by the specific activity of the probe, the rate and extent of the hybridization reaction, the performance of the method for separating hybridized and unhybridized probe, and the sensitivity with which the label can be detected. Under the best conditions, direct hybridization methods such as that described above can detect about  $1 \times 10^5$  to  $1 \times 10^6$  target molecules. The most sensitive procedures may lack many of the features

required for routine clinical and environmental testing such as speed, convenience, and economy. Furthermore, their sensitivities may not be sufficient for many desired applications. Infectious diseases may be associated with  
5 as few as one pathogenic microorganism per 10 ml of blood or other specimen. Forensic investigators may have available only trace amounts of tissue available from a crime scene. Researchers may need to detect and/or quantify a specific gene sequence that is present as only a  
10 tiny fraction of all the sequences present in an organism's genetic material or in the messenger RNA population of a group of cells.

As a result of the interactions among the various components and component steps of this type of assay,  
15 there is almost always an inverse relationship between sensitivity and specificity. Thus, steps taken to increase the sensitivity of the assay (such as increasing the specific activity of the probe) may result in a higher percentage of false positive test results. The linkage  
20 between sensitivity and specificity has been a significant barrier to improving the sensitivity of hybridization assays. One solution to this problem would be to specifically increase the amount of target sequence present using an amplification procedure. Amplification  
25 of a unique portion of the target sequence without requiring amplification of a significant portion of the information encoded in the remaining sequences of the sample could give an increase in sensitivity while at the same time not compromising specificity. For example, a  
30 nucleic acid sequence of 25 bases in length has a probability of occurring by chance of 1 in  $4^{25}$  or 1 in  $10^{15}$  since each of the 25 positions in the sequence may be occupied by one of four different nucleotides.

A method for specifically amplifying nucleic acid  
35 sequences termed the "polymerase chain reaction" or "PCR" has been described by Mullis et al. (See U.S. patents 4,683,195, 4,683,202 and 4,800,159 and European patent

applications 86302298.4, 86302299.2, and 87300203.4 and Methods in Enzymology, Volume 155, 1987, pp. 335-350). The procedure uses repeated cycles of primer-dependent nucleic acid synthesis occurring simultaneously using each  
5 strand of a complementary sequence as a template. The sequence which is amplified is defined by the locations of the primer molecules that initiate synthesis. The primers are complementary to the 3'-terminal portion of the target sequence or its complement and must complex with those  
10 sites in order for nucleic acid synthesis to begin. After extension product synthesis, the strands are separated, generally by thermal denaturation, before the next synthesis step. In the PCR procedure, copies of both strands of a complementary sequence are synthesized.

15 The strand separation step used in PCR to separate the newly synthesized strands at the conclusion of each cycle of the PCR reaction is often thermal denaturation. As a result, either a thermostable enzyme is required or new enzyme must be added between thermal denaturation  
20 steps and the initiation of the next cycle of DNA synthesis. The requirement of repeated cycling of reaction temperature between several different and extreme temperatures is a disadvantage of the PCR procedure. In order to make the PCR convenient, expensive programmable  
25 thermal cycling instruments are required.

The PCR procedure has been coupled to RNA transcription by incorporating a promoter sequence into one of the primers used in the PCR reaction and then, after amplification by the PCR procedure for several  
30 cycles, using the double-stranded DNA as template for the transcription of single-stranded RNA. (See, e.g. Murakawa et al., DNA 7:287-295 (1988).

Other methods for amplification of a specific nucleic acid sequence comprise a series of primer hybridization,  
35 extending and denaturing steps to provide an intermediate double stranded DNA molecule containing a promoter sequence through the use of a primer. The double stranded

DNA is used to produce multiple RNA copies of the target sequence. The resulting RNA copies can be used as target sequences to produce further copies and multiple cycles can be performed. (See, e.g., Burg, et al., WO 89/1050  
5 and Gingeras, et al., WO 88/10315.)

Methods for chemically synthesizing relatively large amounts of DNA of a specified sequence in vitro are well known to those skilled in the art; production of DNA in this way is now commonplace. However, these procedures  
10 are time-consuming and cannot be easily used to synthesize oligonucleotides much greater in length than about 100 bases. Also, the entire base sequence of the DNA to be synthesized must be known. These methods require an expensive instrument capable of synthesizing only a single  
15 sequence at one time. Operation of this instrument requires considerable training and expertise. Methods for the chemical synthesis of RNA have been more difficult to develop.

Nucleic acids may be synthesized by techniques which  
20 involve cloning or insertion of specific nucleic acid sequences into the genetic material of microorganisms so that the inserted sequences are replicated when the organism replicates. If the sequences are inserted next to and downstream from a suitable promoter sequence, RNA  
25 copies of the sequence or protein products encoded by the sequence may be produced. Although cloning allows the production of virtually unlimited amounts of specific nucleic acid sequences, due to the number of manipulations involved it may not be suitable for use in  
30 diagnostic, environmental, or forensic testing. Use of cloning techniques requires considerable training and expertise. The cloning of a single sequence may consume several man-months of effort or more.

Relatively large amounts of certain RNAs may be made  
35 using a recombinant single-stranded-RNA molecule having a recognition sequence for the binding of an RNA-directed polymerase, preferably Q $\beta$  replicase. (See, e.g., U.S.

Patent No. 4,786,600 to Kramer, et al..) A number of steps are required to insert the specific sequence into a DNA copy of the variant molecule, clone it into an expression vector, transcribe it into RNA and then replicate it with  
5 Q $\beta$  replicase.

#### Summary of the Invention

The present invention is directed to novel methods of synthesizing multiple copies of a target nucleic acid  
10 sequence which are autocatalytic (i.e., able to cycle automatically without the need to modify reaction conditions such as temperature, pH, or ionic strength and using the product of one cycle in the next one).

The present method includes (a) treating an RNA  
15 target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target to complex therewith and which optionally has a sequence 5' to the priming sequence which includes a  
20 promoter for an RNA polymerase under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated, (b) extending the first primer in an extension reaction using the target as a template to give a first DNA primer extension product complementary to  
25 the RNA target, (c) separating the DNA extension product from the RNA target using an enzyme which selectively degrades the RNA target; (d) treating the DNA primer extension product with a second oligonucleotide which comprises a primer or a splice template and which has a  
30 complexing sequence sufficiently complementary to the 3'-terminal portion of the DNA primer extension product to complex therewith under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated, provided that if the first  
35 oligonucleotide does not have a promoter, then the second oligonucleotide is a splice template which has a sequence 5' to the complexing sequence which includes a promoter

for an RNA polymerase; (e) extending the 3'-terminus of either the second oligonucleotide or the first primer extension product, or both, in a DNA extension reaction to produce a template for the RNA polymerase; and (f) using the template to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter sequence. The oligonucleotide and RNA copies may be used to autocatalytically synthesize multiple copies of the target sequence.

10 In one aspect of the present invention, the general method includes (a) treating an RNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target to complex there-  
15 with and which has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase under conditions whereby an oligonucleotide/target complex is formed and DNA synthesis may be initiated, (b) extending the first primer in an extension reaction using the  
20 target as a template to give a first DNA primer extension product complementary to the RNA target, (c) separating the first DNA primer extension product from the RNA target using an enzyme which selectively degrades the RNA target; (d) treating the DNA primer extension product with a  
25 second oligonucleotide which comprises a second primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the DNA primer extension product to complex therewith under conditions whereby an oligonucleotide/target complex is formed and DNA synthesis  
30 may be initiated; (e) extending the 3'-terminus of the second primer in a DNA extension reaction to give a second DNA primer extension product, thereby producing a template for the RNA polymerase; and (f) using the template to produce multiple RNA copies of the target sequence using  
35 an RNA polymerase which recognizes the promoter sequence. The oligonucleotide and RNA copies may be used to autocatalytically synthesize multiple copies of the target

sequence. This aspect further includes: (g) treating an RNA copy from step (f) with the second primer under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated; (h) 5 extending the 3' terminus of the second primer in a DNA extension reaction to give a second DNA primer extension product using the RNA copy as a template; (i) separating the second DNA primer extension product from the RNA copy using an enzyme which selectively degrades the RNA copy; 10 (j) treating the second DNA primer extension product with the first primer under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated; (k) extending the 3' terminus of the second primer extension product in a DNA extension 15 reaction to produce a template for an RNA polymerase; and (l) using the template of step (k) to produce multiple copies of the target sequence using an RNA polymerase which recognizes the promoter. Using the RNA copies of step (l), steps (g) to (k) may be autocatalytically 20 repeated to synthesize multiple copies of the target sequence. The first primer which in step (k) acts as a splice template may also be extended in the DNA extension reaction of step (k).

Another aspect of the general method of the present 25 invention provides a method which comprises (a) treating an RNA target sequence with a first primer which has a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex therewith under conditions whereby an 30 oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated; (b) extending the 3' terminus of the primer in an extension reaction using the target as a template to give a DNA primer extension product complementary to the RNA target; (c) separating the DNA 35 extension product from the RNA target using an enzyme which selectively degrades the RNA target; (d) treating the DNA primer extension product with a second

oligonucleotide which comprises a splice template which has a complexing sequence sufficiently complementary to the 3'-terminus of the primer extension product to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated; (e) extending the 3' terminus of the DNA primer extension product to add thereto a sequence complementary to the promoter, thereby producing a template for an RNA polymerase; (f) using the template to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter sequence; and (g) using the RNA copies of step (f), autocatalytically repeating steps (a) to (f) to amplify the target sequence. Optionally, the splice template of step (d) may also function as a primer and in step (e) be extended to give a second primer extension product using the first primer extension product as a template.

20 In addition, in another aspect of the present invention, where the sequence sought to be amplified is present as DNA, use of an appropriate Preliminary Procedure generates RNA copies which may then be amplified according to the General Method of the present invention.

25 Accordingly, in another aspect, the present invention is directed to Preliminary Procedures for use in conjunction with the amplification method of the present invention which not only can increase the number of copies present to be amplified, but also can provide RNA copies of a DNA sequence for amplification.

30 The present invention is directed to methods for increasing the number of copies of a specific target nucleic acid sequence in a sample. In one aspect, the present invention involves cooperative action of a DNA polymerase (such as a reverse transcriptase) and a DNA-dependent RNA polymerase (transcriptase) with an enzymatic hybrid-separation step to produce products that may them-

selves be used to produce additional product, thus resulting in an autocatalytic reaction without requiring manipulation of reaction conditions such as thermal cycling. In some embodiments of the methods of the present invention which include a Preliminary Procedure, all but the initial step(s) of the preliminary procedure are carried out at one temperature.

The methods of the present invention may be used as a component of assays to detect and/or quantitate specific nucleic acid target sequences in clinical, environmental, forensic, and similar samples or to produce large numbers of copies of DNA and/or RNA of specific target sequence for a variety of uses. These methods may also be used to produce multiple DNA copies of a DNA target sequence for cloning or to generate probes or to produce RNA and DNA copies for sequencing.

In one example of a typical assay, a sample to be amplified is mixed with a buffer concentrate containing the buffer, salts, magnesium, nucleotide triphosphates, primers and/or splice templates, dithiothreitol, and spermidine. The reaction is then optionally incubated near 100°C for two minutes to denature any secondary structure. After cooling to room temperature, if the target is a DNA target without a defined 3' terminus, reverse transcriptase is added and the reaction mixture is incubated for 12 minutes at 42°C. The reaction is again denatured near 100°C, this time to separate the primer extension product from the DNA template. After cooling, reverse transcriptase, RNA polymerase, and RNase H are added and the reaction is incubated for two to four hours at 37°C. The reaction can then be assayed by denaturing the product, adding a probe solution, incubating 20 minutes at 60°C, adding a solution to selectively hydrolyze the unhybridized probe, incubating the reaction six minutes at 60°C, and measuring the remaining chemiluminescence in a luminometer. (See, e.g., Arnold, et al., PCT US88/02746 (filed September 21, 1988,

published March 29, 1989) the disclosure of which is incorporated herein by reference and is referred to as "HPA"). The products of the methods of the present invention may be used in many other assay systems known to those skilled in the art.

If the target has a defined 3' terminus or the target is RNA, a typical assay includes mixing the target with the buffer concentrate mentioned above and denaturing any secondary structure. After cooling, reverse transcriptase, RNA polymerase, and RNase H are added and the mixture is incubated for two to four hours at 37°C. The reaction can then be assayed as described above.

The methods of the present invention and the materials used therein may be incorporated as part of diagnostic kits for use in diagnostic procedures.

#### Definitions

As used herein, the following terms have the following meanings unless expressly stated to the contrary.

##### 1. Template

A "template" is a nucleic acid molecule that is being copied by a nucleic acid polymerase. A template may be either single-stranded, double-stranded or partially double-stranded, depending on the polymerase. The synthesized copy is complementary to the template or to at least one strand of a double-stranded or partially double-stranded template. Both RNA and DNA are always synthesized in the 5' to 3' direction and the two strands of a nucleic acid duplex always are aligned so that the 5' ends of the two strands are at opposite ends of the duplex (and, by necessity, so then are the 3' ends).

## 2. Primer, Splice Template

A "primer" is an oligonucleotide that is complementary to a template which complexes (by hydrogen bonding or hybridization) with the template to give a primer/  
5 template complex for initiation of synthesis by a DNA polymerase, and which is extended by the addition of covalently bonded bases linked at its 3' end which are complementary to the template in the process of DNA synthesis. The result is a primer extension product.  
10 Virtually all DNA polymerases (including reverse transcriptases) that are known require complexing of an oligonucleotide to a single-stranded template ("priming") to initiate DNA synthesis, whereas RNA replication and transcription (copying of RNA from DNA) generally do not  
15 require a primer. Under appropriate circumstances, a primer may act as a splice template as well (see definition of "splice template" that follows).

A "splice template" is an oligonucleotide that complexes with a single-stranded nucleic acid and is used  
20 as a template to extend the 3' terminus of a target nucleic acid to add a specific sequence. The splice template is sufficiently complementary to the 3' terminus of the target nucleic acid molecule, which is to be extended, to complex therewith. A DNA- or RNA-dependent  
25 DNA polymerase is then used to extend the target nucleic acid molecule using the sequence 5' to the complementary region of the splice template as a template. The extension product of the extended molecule has the specific sequence at its 3'-terminus which is complementary to the  
30 sequence at the 5'-terminus of the splice template.

If the 3' terminus of the splice template is not blocked and is complementary to the target nucleic acid, it may also act as a primer and be extended by the DNA polymerase using the target nucleic acid molecule as a  
35 template. The 3' terminus of the splice template can be blocked in a variety of ways, including having a 3'-terminal dideoxynucleotide or a 3'-terminal sequence non-

complementary to the target, or in other ways well known to those skilled in the art.

Either a primer or a splice template may complex with a single-stranded nucleic acid and serve a priming  
5 function for a DNA polymerase.

### 3. Target Nucleic Acid, Target Sequence

A "target nucleic acid" has a "target sequence" to be amplified, and may be either single-stranded or double-  
10 stranded and may include other sequences besides the target sequence which may not be amplified.

The term "target sequence" refers to the particular nucleotide sequence of the target nucleic acid which is to be amplified. The "target sequence" includes the  
15 complexing sequences to which the oligonucleotides (primers and/or splice template) complex during the processes of the present invention. Where the target nucleic acid is originally single-stranded, the term "target sequence" will also refer to the sequence comple-  
20 mentary to the "target sequence" as present in the target nucleic acid. Where the "target nucleic acid" is originally double-stranded, the term "target sequence" refers to both the (+) and (-) strands.

### 25 4. Promoter/Promoter Sequence

A "promoter sequence" is a specific nucleic acid sequence that is recognized by a DNA-dependent RNA polymerase ("transcriptase") as a signal to bind to the nucleic acid and begin the transcription of RNA at a  
30 specific site. For binding, such transcriptases generally require DNA which is double-stranded in the portion comprising the promoter sequence and its complement; the template portion (sequence to be transcribed) need not be double-stranded. Individual DNA-dependent RNA polymerases  
35 recognize a variety of different promoter sequences which can vary markedly in their efficiency in promoting transcription. When an RNA polymerase binds to a promoter

sequence to initiate transcription, that promoter sequence is not part of the sequence transcribed. Thus, the RNA transcripts produced thereby will not include that sequence.

5

#### 5. DNA-dependent DNA Polymerase

A "DNA-dependent DNA polymerase" is an enzyme that synthesizes a complementary DNA copy from a DNA template. Examples are DNA polymerase I from E. coli and bacteriophage T7 DNA polymerase. All known DNA-dependent DNA polymerases require a complementary primer to initiate synthesis. It is known that under suitable conditions a DNA-dependent DNA polymerase may synthesize a complementary DNA copy from an RNA template.

15

#### 6. DNA-dependent RNA Polymerase (Transcriptase)

A "DNA-dependent RNA polymerase" or "transcriptase" is an enzyme that synthesizes multiple RNA copies from a double-stranded or partially-double stranded DNA molecule having a (usually double-stranded) promoter sequence. The RNA molecules ("transcripts") are synthesized in the 5' → 3' direction beginning at a specific position just downstream of the promoter. Examples of transcriptases are the DNA-dependent RNA polymerase from E. coli and bacteriophages T7, T3, and SP6.

25

#### 7. RNA-dependent DNA polymerase (Reverse Transcriptase)

An "RNA-dependent DNA polymerase" or "reverse transcriptase" is an enzyme that synthesizes a complementary DNA copy from an RNA template. All known reverse transcriptases also have the ability to make a complementary DNA copy from a DNA template; thus, they are both RNA- and DNA-dependent DNA polymerases. A primer is required to initiate synthesis with both RNA and DNA templates.

35

#### 8. RNase H

An "RNase H" is an enzyme that degrades the RNA portion of an RNA:DNA duplex. RNase H's may be endonucleases or exonucleases. Most reverse transcriptase enzymes normally contain an RNase H activity in addition to their polymerase activity. However, other sources of the RNase H are available without an associated polymerase activity. The degradation may result in separation of RNA from a RNA:DNA complex. Alternatively, the RNase H may simply cut the RNA at various locations such that portions of the RNA melt off or permit enzymes to unwind portions of the RNA.

#### 9. Plus/Minus Strand(s)

Discussions of nucleic acid synthesis are greatly simplified and clarified by adopting terms to name the two complementary strands of a nucleic acid duplex. Traditionally, the strand encoding the sequences used to produce proteins or structural RNAs was designated as the "plus" strand and its complement the "minus" strand. It is now known that in many cases, both strands are functional, and the assignment of the designation "plus" to one and "minus" to the other must then be arbitrary. Nevertheless, the terms are very useful for designating the sequence orientation of nucleic acids and will be employed herein for that purpose.

#### 10. Hybridize, Hybridization

The terms "hybridize" and "hybridization" refer to the formation of complexes between nucleotide sequences which are sufficiently complementary to form complexes via Watson-Crick base pairing. Where a primer (or splice template) "hybridizes" with target (template), such complexes (or hybrids) are sufficiently stable to serve the priming function required by the DNA polymerase to initiate DNA synthesis.

### 11. Primer sequences

The sequences of the primers referred to h r in are set forth below.

#### HBV region 2 primers

- 5 (+): 5' CACCAAATGCCCCTATCTTATCAACACTTCCGG3'  
(-): 5' AATTTAATACGACTCACTATAGGGAGACCCGAGATTGAG  
ATCTTCTGCGAC3'

#### Probe

- (+): 5' GGTCCCCTAGAAGAAGAACTCCCTCG3'

#### 10 HIV region 1 primers

- (+): 5' AATTTAATACGACTCACTATAGGGAGACAAGGGACTTTCC  
GCTGGGGACTTTCC3'  
(-): 5' GTCTAACCAGAGAGACCCAGTACAGGC3'

#### Probe sequence:

- 15 5' GCAGCTGCTTATATGCAGGATCTGAGGG3'

#### HIV region 2 primers

- (+): 5' AATTTAATACGACTCACTATAGGGAGACAAATGGCA  
GTATTCATCCACA3'  
(-): 5' CCCTTCACCTTTCCAGAG3'

#### 20 Probe sequence:

- (-): 5' CTA CTATTCTTTCCCCTGCACTGTACCCC3'

#### HIV region 3 primers

- (+): 5' CTCGACGCAGGACTCGGCTTGCTG3'  
(-): 5' AATTTAATACGACTCACTATAGGGAGACTCCCCCGCTT  
25 AATACTGACGCT3'

#### Probe:

- (+): 5' GACTAGCGGAGGCTAGAAGGAGAGAGATGGG3'

#### HIV region 4 primers

- (+): 5' AATTTAATACGACTCACTATAGGGAGAGACCATCAATGAGGAA  
30 GCTGCAGAATG3'  
(-): 5' CCATCCTATTTGTTCTCTCATCTGGCC3'

#### Probe:

- (-): 5' CTTCCCCTTGGTTCTCTCATCTGGCC3'

#### HIV region 5 primers

- 35 (+): 5' GGCAAATGGTACATCAGGCCATATCACCTAG3'  
(-): 5' AATTTAATACGACTCACTATAGGGAGAGGGGTGGCTCCTT  
CTGATAATGCTG3'

17

Prob :

5'GAAGGCTTTCAGCCCAGAAGTAATACCCATG3'

BCL-2 chromosomal translocation major breakpoint t(14;18)  
primers

5 (-): 5'GAATTAATACGACTCACTATAGGGAGACCTGAGGAGACGGTGACC3'

(+): 5'TATGGTGGTTTGACCTTTAG3'

Probes:

5'GGCTTTCTCATGGCTGTCCTTCAG3'

5'GGTCTTCCTGAAATGCAGTGGTCG3'

10 CML chromosomal translocation t(9;22) primers

(-): 5'GAATTAATACGACTCACTATAGGGAGACTCAGAC

CCTGAGGCTCAAAGTC3'

(+): 5'GGAGCTGCAGATGCTGACCAAC3'

Probe:

15 5'GCAGAGTTCAAAGCCCTTCAGCGG3'

## 12. Specificity

Characteristic of a nucleic acid sequence which describes its ability to distinguish between target and  
20 non-target sequences dependent on sequence and assay conditions.

## Brief Description of the Drawing

FIGs. 1A to 10 depict the General Methods of the  
25 present invention.

FIGs. 2A to 2E depict the embodiment of the present invention referred to as Preliminary Procedure I.

FIG. 3 depicts the embodiment of the present invention referred to as Preliminary Procedure II.

30 FIG. 4A to 4D depicts the improved amplification method.

FIG. 5 shows the results of experiments testing the hypothesis that RNase H from AMV and MMLV and E. coli have specific RNA cleavage sites.

35 FIG. 6 shows the results of incorporation of <sup>32</sup>P-labeled primers during amplification.

Detailed Description of the Invention

In accordance with the present invention, novel methods and compositions are provided for the amplification of specific nucleic acid target sequences for use in assays for the detection and/or quantitation of specific nucleic acid target sequences or for the production of large numbers of copies of DNA and/or RNA of specific target sequences for a variety of uses.

10 I. General Method

In a preferred aspect, the present invention provides an autocatalytic amplification method which synthesizes large numbers of DNA and RNA copies of an RNA target sequence. The target nucleic acid contains the target sequence to be amplified. The target sequence is that region of the target nucleic acid which is defined on either end by the primers, splice templates, and/or the natural target nucleic acid termini and includes both the (+) and (-) strands.

20 In one aspect, this method comprises treating a target nucleic acid comprising an RNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith and which optionally has a sequence 5' to the complexing sequence which includes a promoter sequence for an RNA polymerase under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated. The first oligonucleotide primer may also have other sequences 5' to the priming sequence. The 3'-end of the first primer is extended by an appropriate DNA polymerase in an extension reaction using the RNA as a template to give a first DNA primer extension product which is complementary to the RNA template. The first primer extension product is separated (at least partially) from the RNA template using an enzyme which selectively degrades the RNA template. Suitable

enzymes are those which selectively act on the RNA strand of an RNA-DNA complex and include enzymes which comprise an RNase H. Although some reverse transcriptases include an RNase H activity, it may be preferable to add exogenous RNase H, such as an E. coli RNase H.

The single-stranded first primer extension product is treated with a second oligonucleotide which comprises a second primer or a splice template which has a complexing sequence sufficiently complementary to the 3'-terminal portion of target sequence contained in the first primer extension product to complex therewith, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated. If the first primer does not have a promoter then the second oligonucleotide is a splice template which has a sequence 5' to the complexing region which includes a promoter for an RNA polymerase. Optionally, the splice template may be blocked at its 3' terminus. The 3' terminus of the second oligonucleotide and/or the primer extension product is extended in a DNA extension reaction to produce a template for a RNA polymerase. The RNA copies or transcripts produced may autocatalytically multiply without further manipulation.

Where an oligonucleotide functions as a splice template, its primer function is not required. Thus, the 3' terminus of the splice template may be either blocked or unblocked. The components of the resulting reaction mixture (i.e., an RNA target which allows production of a first primer extension product with a defined 3' terminus, a first primer, and a splice template either blocked or unblocked) function to autocatalytically synthesize large quantities of RNA and DNA.

In one aspect of the present invention, the first and second oligomers both are primers. The first primer has a sequence 5' to the complexing sequence which includes a promoter for a RNA polymerase and may include other sequences. The second primer may also include a sequence

5' to the complexing sequence which may include a promoter for an RNA polymerase and optionally other sequences. Where both primers have a promoter sequence, it is preferred that both sequences are recognized by the same RNA polymerase unless it is intended to introduce the second promoter for other purposes, such as cloning. The 3'-end of the second primer is extended by an appropriate DNA polymerase in an extension reaction to produce a second DNA primer extension product complementary to the first primer extension product. Note that as the first primer defined one end of the target sequence, the second primer now defines the other end. The double-stranded product of the second extension reaction is a suitable template for the production of RNA by an RNA polymerase. If the second primer also has a promoter sequence, transcripts complementary to both strands of the double-stranded template will be produced during the autocatalytic reaction. The RNA transcripts may now have different termini than the target nucleic acid, but the sequence between the first primer and the second primer remains intact. The RNA transcripts so produced may automatically recycle in the above system without further manipulation. Thus, this reaction is autocatalytic.

If the complexing sequence of the second primer complexes with the 3' terminus of the first primer extension product, the second primer may act as a splice template and the first primer extension product may be extended to add any sequence of the second primer 5' to the priming sequence to the first primer extension product. (See, e.g., Figures 1E and 1G) If the second primer acts as a splice template and includes a promoter sequence 5' to the complexing sequence, extension of the first primer extension product to add the promoter sequence produces an additional template for an RNA polymerase which may be transcribed to produce RNA copies of either strand. (See Figures 1E and 1G) Inclusion of

promoters in both primers may enhance the number of copies of the target sequence synthesized.

Another aspect of the general method of the present invention includes using a first oligonucleotide which  
5 comprises a primer and a second oligonucleotide which comprises a splice template and which may or may not be capable of acting as a primer per se (in that it is not itself extended in a primer extension reaction). This aspect of the general method comprises treating a target  
10 nucleic acid comprising an RNA target sequence with a first oligonucleotide primer which has a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex therewith under conditions whereby an oligonucleotide/target sequence  
15 complex is formed and DNA synthesis may be initiated. The first primer may have other sequences 5' to the complexing sequence, including a promoter. The 3' end of the first primer is extended by an appropriate DNA polymerase in an extension reaction using the RNA as a template to give a  
20 first primer extension product which is complementary to the RNA template. The first primer extension product is separated from the RNA template using an enzyme which selectively degrades the RNA template. Suitable enzymes are those which selectively act on the RNA strand of an  
25 RNA-DNA complex and include enzymes which comprise an RNase H activity. Although some reverse transcriptases include an RNase H activity, it may be preferable to add exogenous RNase H, such as an E. coli RNase H. The single stranded first primer extension product is treated with a  
30 splice template which has a complexing sequence sufficiently complementary to the 3'-terminus of the primer extension product to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase under conditions whereby an  
35 oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated. The 3' terminus of the splice template may be either blocked (such as by addition of a

dideoxynucleotide) or uncomplementary to the target nucleic acid (so that it does not function as a primer) or alternatively unblocked. The 3' terminus of the first primer extension product is extended using an appropriate  
5 DNA polymerase in a DNA extension reaction to add to the 3' terminus of the first primer extension product a sequence complementary to the sequence of the splice template 5' to the complexing sequence which includes the promoter. If the 3' terminus is unblocked, the splice  
10 template may be extended to give a second primer extension product complementary to the first primer extension product. The product of the extension reaction with the splice template (whether blocked or unblocked) can function as a template for RNA synthesis using an RNA  
15 polymerase which recognizes the promoter. As noted above, RNA transcripts so produced may automatically recycle in the above system without further manipulation. Thus, the reaction is autocatalytic.

In some embodiments, the target sequence to be  
20 amplified is defined at both ends by the location of specific sequences complementary to the primers (or splice templates) employed. In other embodiments, the target sequence is defined at one location of a specific sequence, complementary to a primer molecule employed and,  
25 at the opposite end, by the location of a specific sequence that is cut by a specific restriction endonuclease, or by other suitable means, which may include a natural 3' terminus. In other embodiments, the target sequence is defined at both ends by the location of  
30 specific sequences that are cut by one or more specific restriction endonuclease(s).

In a preferred embodiment of the present invention, the RNA target sequence is determined and then analyzed to determine where RNase H degradation will cause cuts or  
35 removal of sections of RNA from the duplex. Analyses can be conducted to determine the effect of the RNase degradation of the target sequence by RNase H present in

AMV reverse transcriptase and MMLV reverse transcriptase, by E. coli RNase H or other sources and by combinations thereof.

In selecting a primer set, it is preferable that one  
5 of the primers be selected so that it will hybridize to a section of RNA which is substantially nondegraded by the RNase H present in the reaction mixture. If there is substantial degradation, the cuts in the RNA strand in the region of the primer may inhibit initiation of DNA  
10 synthesis and prevent extension of the primer. Thus, it is preferred to select a primer which will hybridize with a sequence of the RNA target, located so that when the RNA is subjected to RNase H, there is no substantial degradation which would prevent formation of the primer  
15 extension product.

The site for hybridization of the promoter-primer is chosen so that sufficient degradation of the RNA strand occurs to permit removal of the portion of the RNA strand hybridized to the portion of the DNA strand to which the  
20 promoter-primer will hybridize. Typically, only portions of RNA are removed from the RNA:DNA duplex through RNase H degradation and a substantial part of the RNA strand remains in the duplex.

Formation of the promoter-containing double stranded  
25 product for RNA synthesis is illustrated in Figure 4. As illustrated in Figure 4, the target RNA strand hybridizes to a primer which is selected to hybridize with a region of the RNA strand which is not substantially degraded by RNase H present in the reaction mixture. The primer is  
30 then extended to form a DNA strand complementary to the RNA strand. Thereafter, the RNA strand is cut or degraded at various locations by the RNase H present in the reaction mixture. It is to be understood that this cutting or degradation can occur at this point or at other  
35 times during the course of the reaction. Then the RNA fragments dissociate from the DNA strand in regions where

significant cuts or degradation occur. The promoter-primer then hybridizes to the DNA strand at its 3' end, where the RNA strand has been substantially degraded and separated from the DNA strand. Next, the DNA strand is  
5 extended to form a double strand DNA promoter sequence, thus forming a template for RNA synthesis. It can be seen that this template contains a double-stranded DNA promoter sequence. When this template is treated with RNA polymerase, multiple strands of RNA are formed.

10 Although the exact nature of the RNA degradation resulting from the RNase H is not known, it has been shown that the result of RNase H degradation on the RNA strand of an RNA:DNA hybrid resulted in dissociation of small pieces of RNA from the hybrid. It has also been shown  
15 that promoter-primers can be selected which will bind to the DNA after RNase H degradation at the area where the small fragments are removed.

Figures 1 and 2, as drawn, do not show the RNA which may remain after RNase H degradation. It is to be  
20 understood that although these figures generally show complete removal of RNA from the DNA:RNA duplex, under the preferred conditions only partial removal occurs as illustrated in Figure 3. By reference to Figure 1A, it can be seen that the proposed mechanism may not occur if  
25 a substantial portion of the RNA strand of Figure 1 remains undegraded thus preventing hybridization of the second primer or extension of the hybridized second primer to produce a DNA strand complementary to the promoter sequence. However, based upon the principles of synthesis  
30 discovered and disclosed in this application, routine modifications can be made by those skilled in the art according to the teachings of this invention to provide an effective and efficient procedure for amplification of RNA.

35 As may be seen from the descriptions herein and Figures 1A to 10, the method of the present invention embraces optimal variations.

Figure 1A depicts a method according to the present invention wherein the target nucleic acid has additional sequences 3' to the target sequence. The first oligonucleotide comprises a first primer having a promoter 5' to its complexing sequence which complexes with the 3' terminal portion of the target sequence of a target nucleic acid (RNA) which has additional sequences 3' to the end of the target sequence. The second oligonucleotide comprises a second primer which complexes with the 3' terminal portion of the first primer extension product, coinciding with the 3' terminus of the first primer extension product. In step (1), the first primer does not act as a splice template due to the additional sequences 3' to the target sequence; however, in step (10), the first primer can act as a splice template, since the second primer extension product does not have additional sequences 3' to the target sequence.

Figure 1B depicts a method according to the present invention wherein the target nucleic acid (RNA) has additional sequences both 5' and 3' to the target sequence. The first oligonucleotide is as depicted in Figure 1A. The second oligonucleotide comprises a primer which complexes to the 3' terminal portion of the target sequence of the first primer extension product which has additional sequences 3' to the target sequence.

Figure 1C depicts a target nucleic acid (RNA) which has defined 5' and 3' ends and, thus, has no additional sequences either 5' or 3' to the target sequence. The first oligonucleotide is as depicted in Figures 1A and 1B, but since it complexes with the 3' terminus of the target nucleic acid, it acts as both a primer and splice template in Step 1. The second oligonucleotide is as depicted in Figure 1A.

Figure 1D depicts a target nucleic acid (RNA) having a defined 3' end and, thus, has no additional sequences 3' to the target sequence, but does have additional sequences 5' to the target sequence. The first oligonucleotide is

as depicted in Figure 1C and functions as both a primer and a splice template. The second oligonucleotide is as depicted in Figure 1B.

Figure 1E depicts a target nucleic acid (RNA) which has a defined 5' end but which has additional sequences 3' to the target sequence. The first oligonucleotide is as depicted in Figure 1A. The second oligonucleotide comprises a second primer which, since it complexes with the 3'-terminus of the first primer extension product, also comprises a splice template. The second oligonucleotide also has a promoter 5' to its complexing sequence.

Figure 1F depicts a target nucleic acid (RNA) having additional sequences both 5' and 3' to the target sequence. The first oligonucleotide is as depicted in Figure 1A. The second oligonucleotide is as depicted in Figure 1E, except it cannot act as a splice template in step (4), since the first primer extension product has additional sequences 3' to the target sequence.

Figure 1G depicts a target nucleic acid (RNA) which has both defined 5' and 3' ends, having no sequences besides the target sequence. The first oligonucleotide is as depicted in Figure 1C and the second oligonucleotide as depicted in Figure 1E. Since the target has no additional sequences, both oligonucleotides also act as splice templates.

Figure 1H depicts a target nucleic acid (RNA) which has a defined 3' end, having no sequences 3' to the target sequence, but has additional sequences 5' to the target sequence. The first oligonucleotide is as depicted in Figures 1C and 1G and acts as both a primer and splice template. The second oligonucleotide is as depicted in Figure 1F.

Figure 1I depicts a target nucleic acid (RNA) which has a defined 5' terminus and has no additional sequences 5' to the target sequence, but has additional sequences 3' to the target sequence. The first oligonucleotide

comprises a primer without a promoter. The second oligonucleotide comprises an unblocked splice template which has a promoter 5' to its complexing sequence.

Figure 1J depicts a target nucleic acid (RNA) which  
5 has defined 5' and 3' terminus and no sequences besides the target sequence. The first oligonucleotide comprises a primer without a promoter. The second oligonucleotide is as depicted in Figure 1I.

Figure 1K depicts a target nucleic acid (RNA) which  
10 has a defined 5' terminus with no additional sequence 5' to the target sequence, but which has additional sequences 3' to the target sequence. The second oligonucleotide comprises a splice template having a promoter 5' to its complexing sequence, but which is blocked at its 3'  
15 terminus. The second oligonucleotide is incapable of acting as a primer.

Figure 1L depicts a target nucleic acid (RNA) which has defined 5' and 3' ends and no additional sequences besides the target sequence. The first oligonucleotide  
20 acts as both a primer and a splice template. The second oligonucleotide is a blocked splice template and is as depicted in Figure 1K.

Figure 1M depicts a target nucleic acid (RNA) which has a defined 5'-terminus, and, thus, no additional  
25 sequences 5' to the target sequence, but which has additional sequences 3' to the target sequence. The first

Figure 10 depicts a target nucleic acid (RNA) which has a defined 5' terminus, having no additional sequences 5' to the target sequence, but which has additional sequences 3' to the target sequence. One oligonucleotide is used for both the first and second oligonucleotide. The oligonucleotide has a 3'-primer sequence which complexes to the 3'-terminal portion of the target sequence as shown in Step (1), and has a 5' splice template sequence with a promoter which complexes with the 3' terminus of the primer extension product as shown in step (4).

In summary, the methods of the present invention provide a method for autocatalytically synthesizing multiple copies of a target nucleic acid sequence without repetitive manipulation of reaction conditions such as temperature, ionic strength and pH which comprises (a) combining into a reaction mixture a target nucleic acid which comprises an RNA target sequence; two oligonucleotide primers, a first oligonucleotide having a complexing sequence sufficiently complementary to the 3' terminal portion of the RNA target sequence (for example the (+) strand) to complex therewith and a second oligonucleotide having a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence of its complement (for example, the (-) strand) to complex therewith, wherein the first oligonucleotide comprises a first primer which optionally has a sequence 5' to the complexing sequence which includes a promoter and the second oligonucleotide comprises a primer or a splice template; provided that if the first oligonucleotide does not have a promoter, then the second oligonucleotide is a splice template which has a sequence 5' to the priming sequence which includes a promoter for an RNA polymerase; a reverse transcriptase or RNA and DNA dependent DNA polymerases; an enzyme activity which selectively degrades the RNA strand of an RNA-DNA complex (such as an RNase H) and an RNA polymerase which recognizes the

promoter. The components of the reaction mixture may be combined stepwise or at once. The reaction mixture is incubated under conditions whereby an oligonucleotide/-target sequence is formed, including DNA priming and  
5 nucleic acid synthesizing conditions (including ribonucleotide triphosphates and deoxyribonucleotide triphosphates) for a period of time sufficient whereby multiple copies of the target sequence are produced. The reaction advantageously takes place under conditions suitable for  
10 maintaining the stability of reaction components such as the component enzymes and without requiring modification or manipulation of reaction conditions during the course of the amplification reaction. Accordingly, the reaction may take place under conditions that are substantially  
15 isothermal and include substantially constant ionic strength and pH.

The present reaction does not require a denaturation step to separate the RNA-DNA complex produced by the first DNA extension reaction. Such steps require manipulation  
20 of reaction conditions such as by substantially increasing the temperature of the reaction mixture (generally from ambient temperature to about 80°C to about 105°C), reducing its ionic strength (generally by 10X or more) or changing pH (usually increasing pH to 10 or more). Such  
25 manipulations of the reaction conditions often deleteriously affect enzyme activities, requiring addition of additional enzyme and also necessitate further manipulations of the reaction mixture to return it to conditions suitable for further nucleic acid synthesis.

30 Suitable DNA polymerases include reverse transcriptases. Particularly suitable DNA polymerases include AMV reverse transcriptase and MMLV reverse transcriptase.

Promoters or promoter sequences suitable for incorporation in the primers and/or splice templates used  
35 in the methods of the present invention are nucleic acid sequences (either naturally occurring, produced synthetically or a product of a restriction digest) that

are specifically recognized by an RNA polymerase that recognizes and binds to that sequence and initiates the process of transcription whereby RNA transcripts are produced. The sequence may optionally include nucleotide  
5 bases extending beyond the actual recognition site for the RNA polymerase which may impart added stability or susceptibility to degradation processes or increased transcription efficiency. Promoter sequences for which there is a known and available polymerase that is capable  
10 of recognizing the initiation sequence are particularly suitable to be employed. Typical, known and useful promoters include those which are recognized by certain bacteriophage polymerases such as those from bacteriophage T3, T7 or SP6, or a promoter from E. coli.

15 Although some of the reverse transcriptases suitable for use in the methods of the present invention have an RNase H activity, such as AMV reverse transcriptase, it may be preferred to add exogenous RNase H, such as E. coli RNase H. Although, as the examples show, the addition of  
20 exogenous RNase H is not required, under certain conditions, the RNase H activity present in AMV reverse transcriptase may be inhibited by components present in the reaction mixture. In such situations, addition of exogenous RNase H may be desirable. Where relatively  
25 large amounts of heterologous DNA are present in the reaction mixture, the native RNase H activity of the AMV reverse transcriptase may be somewhat inhibited (see e.g., Example 8) and thus the number of copies of the target sequence produced accordingly reduced. In situations  
30 where the target sequence comprises only a small portion of DNA present (e.g., where the sample contains significant amounts of heterologous DNA), it is particularly preferred to add exogenous RNase H. One such preferred RNase H is E. coli RNase H. Addition of such exogenous  
35 RNase H has been shown to overcome inhibition caused by large amounts of DNA. (Se , e.g., Example 8).

The RNA transcripts produced by these methods may serve as templates to produce additional copies of the target sequence through the above-described mechanisms. The system is autocatalytic and amplification by the methods of the present invention occurs autocatalytically without the need for repeatedly modifying or changing reaction conditions such as temperature, pH, ionic strength or the like. This method does not require an expensive thermal cycling apparatus, nor does it require several additions of enzymes or other reagents during the course of an amplification reaction.

The methods of the present invention may be used as a component of assays to detect and/or quantitate specific nucleic acid target sequences in clinical, environmental, forensic, and similar samples or to produce large numbers of copies of DNA and/or RNA of specific target sequence for a variety of uses.

In a typical assay, a sample to be amplified is mixed with a buffer concentrate containing the buffer, salts, magnesium, triphosphates, primers and/or splice templates, dithiothreitol, and spermidine. The reaction may optionally be incubated near 100°C for two minutes to denature any secondary structures in the nucleic acid. After cooling, if the target is a DNA target without a defined 3' terminus, reverse transcriptase is added and the reaction mixture is incubated for 12 minutes at about 42°C. The reaction is again denatured near 100°C, this time to separate the primer extension product from the DNA template. After cooling, reverse transcriptase, RNA polymerase, and RNase H are added and the reaction is incubated for two to four hours at 37°C. The reaction can then be assayed by denaturing the product, adding a probe solution, incubating 20 minutes at 60°C, adding a solution to selectively hydrolyze the label of the unhybridized probe, incubating the reaction six minutes at 60°C, and measuring the remaining chemiluminescent label in a luminometer. The methods of the HPA application cited

supra are incorporated herein. Several other methods for product determination may be employed in place of the in-solution probe hybridization.

If the target has a defined 3' terminus and one of the oligonucleotides is a splice template which has a complexing sequence sufficiently complementary to the 3' terminus to complex therewith and a promoter sequence 5' to the complex sequence or the target is RNA, a typical assay includes mixing the target with the buffer concentrate mentioned above and denaturing any secondary structure. After cooling, reverse transcriptase, RNA polymerase, and if desired, RNase H are added and the mixture is incubated for two to four hours at 37°C. The reaction can then be assayed as described above.

15

## II. Preliminary Procedures

The following are several embodiments of preliminary procedures which optionally may be employed in conjunction with the preferred method of the present invention. Since some target nucleic acids require modification prior to autocatalytic amplification, these procedures may be employed to accomplish the modifications. Where the target nucleic acid (and target sequence) is originally DNA, these procedures may be employed to produce RNA copies of the target sequence for use in the General Method. It should be appreciated that these Preliminary Procedures may themselves be repeated and therefore may be used as amplification methods in their own right.

### 30 Preliminary Procedure I

This method gives RNA copies of a target sequence of a target nucleic acid which comprises a (single-stranded) DNA with a defined 3' terminus. Preliminary Procedure I uses two nucleic acid components: a target nucleic acid molecule and an oligonucleotide splice template. This procedure requires a DNA target nucleic acid having a defined 3'-end. If the native 3' terminus is not known or

is unsatisfactory for any reason, a new defined 3' terminus may be created by use of a restriction nuclease, ligation to another sequence, or some other means.

In the following description, (see Figs. 2A to 2C)  
5 the target nucleic acid will arbitrarily have the "minus" sense. Thus, the splice template will have the "plus" sense so as to be sufficiently complementary to the target to complex therewith. The splice template has a complexing sequence sufficiently complementary to the 3'  
10 terminus of the target to complex therewith. The splice template also has a sequence 5' to the complexing sequence which includes a promoter sequence for an RNA polymerase. The splice template may optionally have other sequences 5' to the promoter, between the promoter and complexing  
15 sequences, and/or 3' to the complexing sequence. The splice template may also be modified at the 3' terminus to be "blocked" so that it cannot be extended by adding additional nucleotides in an extension reaction and is rendered incapable of acting as a primer in addition to  
20 acting as a splice template.

Preliminary Procedure I uses two enzyme activities: a DNA-dependent DNA polymerase and a DNA-dependent RNA polymerase.

The target nucleic acid is treated with the splice  
25 template under conditions wherein an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated. In a DNA extension reaction using an appropriate DNA polymerase, a sequence complementary to the sequence of the splice template 5' to  
30 the complexing sequence is added to the 3' terminus of the target DNA. The splice template, if not blocked at the 3' terminus, may also serve as a primer for the DNA polymerase and be extended to give a primer extension product. The product of the extension reaction, either double-  
35 strand or partially double-stranded, target/splice template complex acts as a template for the synthesis RNA transcripts using an RNA polymerase which recognizes the

promoter. The RNA transcripts may be then used for the general method or be used to generate DNA copies of the RNA as follows:

An RNA transcript comprising the target sequence  
5 (having the "plus" sense) is treated with a primer (which nominally has the "minus" sense) which has a complexing sequence sufficiently complementary to the 3' end of the target sequence of the RNA transcript to complex therewith under conditions whereby an oligonucleotide/target  
10 sequence complex is formed and DNA synthesis may be initiated. The primer is then extended in a DNA extension reaction using the RNA transcript as template to generate a DNA primer extension product having the target sequence. The DNA target sequence is separated from the RNA  
15 transcript by either denaturation or by degradation of the RNA and beginning with the splice template, the cycle is repeated. Optionally, the primer may also have additional sequences 5' to the priming sequence. The splice template may also have additional sequences 3' to the complexing  
20 sequence.

In one embodiment, the above method may be practiced using one oligonucleotide by using an oligonucleotide having a sequence which would comprise the primer 3' to the sequence which would comprise the splice template.  
25 (See, e.g. Fig. 2C).

Preliminary Procedure I is further described by reference to Figures 2A to 2E. Figure 2A depicts a target nucleic acid (DNA) which has a defined 3' terminus, having no additional sequences 3' to the target sequence. The  
30 first oligonucleotide comprises both a primer and a splice template and has a promoter 5' to its complexing sequence.

Figure 2B depicts a target nucleic acid (DNA) as shown in Figure 2A. The first oligonucleotide comprises a splice template which is blocked at its 3' end and is  
35 thus incapable of acting as a primer.

Figure 2C depicts a target DNA as shown in Figures 2A and 2B. Figure 2C depicts the use of one oligonucleotide

which has a splice template (with a promoter) sequence 5' to a primer sequence at its 3' end. Thus, the oligonucleotide acts as a blocked splice template in steps (1) and (7) and as a primer (and splice template) in step 5 (4).

Figure 2D depicts a target nucleic acid (DNA) which has a defined 5'-end and additional sequences 3' to the target sequence which undergoes prefatory complexing, primer extending and separating steps (steps 1 and 2) to generate a complementary DNA target having a defined 3'-terminus. The oligonucleotide of steps (1) and (7) comprises a primer which complexes with the 3' terminal portion of the target sequence of the original target DNA (here nominally (+)). The other oligonucleotide (of steps 15 (4) and (10)) comprises an unblocked splice template which complexes with the 3'-end of the complement of the original target.

Figure 2E depicts a target nucleic acid (DNA) which has a defined 5' terminus and additional sequences 3' to the target sequence which undergoes prefatory complexing, extending and separating steps (steps (1) and (2)) to generate a complementary DNA target having a defined 3'-terminus. The oligonucleotide of steps (1) and (7) comprises a primer which complexes with the 3' terminal portion of the target sequence of the original target DNA (here nominally (+)). The oligonucleotide of steps (4) and (10) comprises a blocked splice template which complexes with the 3' terminal portion of the target sequence of the complement of the original target.

The splice template is complexed with the 3' terminus of the target nucleic acid under complexing conditions so that a target/splice template complex is formed and DNA synthesis may be initiated. (step 1) A suitable DNA polymerase is used to extend the 3' terminus of the target nucleic acid to add a sequence complementary to the sequence of the splice template 5' to the complexing sequence. If the 3' terminus of the splice template has

not been blocked, and is sufficiently complementary to the target nucleic acid the splice template may act as a primer so that the 3' terminus of the splice template may also be extended. (Figure 2A) The resulting  
5 target/splice template complex may be either partially or completely double-stranded. (See Figure 2A versus Figure 2B and 2C) At minimum the double-stranded region comprises the promoter sequence and the priming sequence which complexed with the target nucleic acid.

10 The template of Step 2 is transcribed by an appropriate RNA polymerase. The RNA polymerase produces about 5-1000 copies of RNA for each template. In Figures 2A to 2C, the RNA produced is nominally of the "plus" sense and includes the sequence from the 3' end of the  
15 promoter to the 5' end of the target nucleic acid.

The RNA product of Step 3 may be used according to the general method to autocatalytically amplify the target sequence or alternatively may be treated under complexing and priming conditions with a primer which has a  
20 complexing sequence at its 3' terminus sufficiently complementary to the 3' terminal portion of the target sequence in the RNA to complex therewith and optionally includes other sequences 5' to the complexing sequence. The primer is extended using the RNA as a template by an  
25 RNA-dependent DNA polymerase in a primer extension reaction. This produces a product which is at least partially double-stranded and which must be made at least partially single-stranded for further synthesis by degradation of the RNA portion, such as by an RNase H  
30 (step 5) or by some other method such as denaturation.

The DNA produced in Step 6 may be used as a target molecule for Step 1 and treated with a splice template as described above to produce more RNA and DNA. These steps may be cycled to produce any desired level of  
35 amplification. It should also be noted that, by appropriate choice of the splice template and primer(s), this new target molecule (Step 6) may have different

termini than the original molecule. Furthermore, if the primer extension product from the RNA has a promoter sequence 5' to the complexing sequence, a second primer having a complexing sequence of the "plus" sense and  
5 optionally other sequences 5' to the complexing sequence may be used to copy the extended primer product to produce a double-stranded template for an RNA polymerase. A template so produced enables the RNA polymerase to make "minus" sense RNA which may be amplified using the general  
10 method or further amplified using the procedures herein.

#### Preliminary Procedure II

Preliminary Procedure II differs from Preliminary Procedure I in the way the autocatalytic species is generated. The DNA target nucleic acid need not have a defined  
15 3' terminus. In one aspect, a primer containing a promoter sequence 5' to the complexing sequence is used instead of a splice template to introduce the promoter sequence into the template for the RNA polymerase. The  
20 primer has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith and a sequence which includes a promoter for an RNA polymerase 5' to the complexing sequence. The primer is extended with a DNA  
25 polymerase to give a primer extension product.

After separation of the strands, usually by thermal denaturation, a second oligonucleotide of the same sense as the target molecule is used as a primer to synthesize a DNA complement to the first primer extension product.  
30 The second primer may have additional sequences 5' to the complexing region which may include a promoter. Inclusion of a promoter sequence in the second primer may enhance amplification. The second primer may also be a splice template.

35 Preliminary Procedure II is further described with reference to Figure 3. Figure 3 depicts a target DNA which has additional sequences both 5' and 3' to the

target sequence. The first primer has a complexing sequence at its 3' terminus and a sequence 5' to the complexing sequence which includes a promoter sequence and complexes sufficiently with the target to serve a priming function and initiate DNA synthesis. An appropriate DNA polymerase extends the primer to give a primer extension product. The strands are separated by denaturation or other means to yield to a single-stranded promoter containing primer extension product.

10 A second primer is used which has a complexing sequence at its 3' terminus sufficiently complementary to the 3' terminal portion of the target sequence of the first primer extension product and, optionally, other sequences 5' to the complexing sequence which may include  
15 a promoter sequence. The second primer is complexed sufficiently with the primer extension product from Step 3 to serve a priming function and initiate DNA synthesis. The DNA polymerase extends the second primer to give a second primer extension product. The resulting double-  
20 stranded DNA molecule may now serve as a template for the RNA polymerase to generate RNA transcripts for the General Method. As depicted in Figure 3, the RNA molecules produced are nominally of the "plus" sense and may be multiplied using the general method of the present  
25 invention.

Where the complexing sequence of the second primer is complementary to the 3' terminus of the first primer extension product from Step 3 and the second primer includes a promoter sequence 5' to the complexing  
30 sequence, the second primer may serve as a splice template so that the 3'-terminus of the first primer extension product from Step 3 may be further extended to add the promoter sequence and produce a template for the RNA polymerase which produces RNA transcripts of both senses.  
35 The RNA molecules so produced may be amplified using the general method.

In another aspect of the present invention, the second primer acts as a splice template and has a promoter 5' to the complexing sequence, so that the first primer need not have a promoter. In that case, the first primer extension product from Step 2 is further extended to produce a complement to the promoter sequence, thus generating a template for the production of "minus" sense RNA by the RNA polymerase.

By repeating the steps described above, additional RNA and DNA copies of the target sequence may be produced.

### Examples

#### Preface

The following examples of the procedures previously described demonstrate the mechanism and utility of the methods of the present invention. They are not limiting to the inventions and should not be considered as such.

Many of the materials used in one or more examples are similar. To simplify the descriptions in the examples, some of the materials will be abbreviated and described here.

The template referred to as "frag 1" is a double-stranded segment of DNA homologous to a region of DNA from the hepatitis B genome. It has been excised from a plasmid via restriction nuclease digestion and purified from the remaining nucleic acid by chromatographic methods. Subsequent to purification, the fragment has been cut with a restriction endonuclease and purified via phenol extraction and ethanol precipitation to yield the desired target.

The template referred to as "M13L(-)" is a purified single-stranded DNA target containing, as a small fraction of the total sequence, the minus strand target sequence.

Several different primers and splice templates were used in the examples described herein. The olig nucleotide referred to as T7pro(+) contains, near the

5' terminus, a T7 RNA polymerase promoter and, near the 3' terminus, a sequence complementary to the 3' terminus of the minus-strand target sequence to the two templates described above; T7pro(+) also contains other sequence information to enhance performance. The sequence for T7pro(+) is 5'-AATTT AATAC GACTC ACTAT AGGGA GAGGT TATCG CTGGA TGTGT CTGCG GCGT3'.

Another oligonucleotide similar to T7pro(+) is ddT7pro(+). It differs from T7pro(+) in that the 3' terminus has been extended with a dideoxy nucleotide using terminal deoxynucleotidyl transferase. Unlike T7pro(+), ddT7pro(+) is incapable of serving as a primer for DNA synthesis by reverse transcriptase but can act as a splice template.

HBV(-)Pr is a primer which will hybridize to the plus strand of the frag 1 template and is homologous to a sequence within the M13L(-). [HBV(-)Pr is complementary to a sequence 3' to the plus strand sequence homologous to the T7pro(+).] The sequence for HBV(-)Pr is 5'-GAGGA CAAAC GGGCA ACATA CCTTG-3'.

Another oligonucleotide containing a promoter region is T7pro(-). This promoter-primer contains a sequence identical to T7pro(+) but replaces the sequence complementary to the minus target with a sequence complementary to the plus target. The 3' terminus of T7pro(-) is complementary to the 3' terminus of the plus strand of frag 1. The sequence for T7pro(-) is 5'-AATTT AATAC GACTC ACTAT AGGGA GATCC TGGAA TTAGA GGACA AACGG GC-3'. Like the ddT7pro(+), ddT7pro(-) is a 3' blocked oligonucleotide made by extending the 3' terminus with a dideoxynucleotide using terminal deoxynucleotidyl transferase. The ddT7pro(-) cannot serve as a primer but is otherwise similar to T7pro(-).

The templates used in these examples contain substantial sequence between the regions homologous or complementary to the primers and splice templates described above. As the sequence between the

oligonucleotides will be amplified as a result of the invention, quantification of this sequence provides a specific means of measuring the amplification. It has been convenient to assay the products by hybridization techniques using DNA probes specific for the sequences coded for between the oligonucleotide primers and splice templates. Two probes are used in the examples presented below: Probe(+) and Probe(-). Probe(+) is complementary to the minus sense product and Probe(-) is complementary to the plus sense product. The sequence for Probe(+) is 5'-CCTCT TCATC CTGCT GCTAT GCCTC-3' and the sequence for Probe(-) is 5'-GAGC ATAGC AGCAG GATGA AGAGG-3'. The probes used herein have been labeled with a chemiluminescent tag. In the assay, the label on hybridized probe emits light which is measured in a luminometer.

In the following examples, relative amplification was measured as follows. A sample of the amplification reaction mixture (usually 10  $\mu$ l) was diluted to 50  $\mu$ l with 10mM Tris-HCl, pH 8.3, and denatured two minutes at 95°C. After cooling on ice, 50  $\mu$ l of a probe solution containing approximately 75 fmol Probe(+) or Probe(-), 0.2 M lithium succinate, pH 5.2, 21% (w/v) lithium lauryl sulfate, 2 mM EDTA, and 2 mM EGTA, was added to the sample and mixed. The reactions were then incubated 20 minutes at 60°C and cooled. To each hybridization reaction was added 500  $\mu$ l of a solution prepared by adjusting a saturated sodium borate solution to pH 8.5 with hydrochloric acid, then diluting to bring the borate concentration to 0.8 M final and adding Triton X-100 to 5% (v/v) final. The reactions were then mixed and incubated six minutes at 60°C to destroy the chemiluminescent label of the unhybridized probe. This method of destruction of the chemiluminescent label of unhybridized probe is quite specific; only a very small fraction of the unhybridized probe remains chemiluminescent. The reactions were cooled and the remaining chemiluminescence was quantified in a

luminometer upon the addition of 200  $\mu$ l of 1.5 M sodium hydroxide, 0.1% (v/v) hydrogen peroxide. In th assay, hybridized probe emits light which is measured in a luminometer. Since the reaction which destroys the  
5 chemiluminescent label of unhybridized probe is not 100% effective, there is generally a background level of signal present in the range of about 300 to 1300 relative light units (RLU).

Many other assay methods are also applicable,  
10 including assays employing hybridization to isotopically labeled probes, blotting techniques and electrophoresis.

The enzymes used in the following examples are avian myeloblastosis virus reverse transcriptase from Seikagaku America, Inc., T7 RNA polymerase from New England Biolabs  
15 or Epicentre, and Moloney murine leukemia virus (MMLV) reverse transcriptase and E. coli RNase H from Bethesda Research Laboratories. Other enzymes containing similar activities and enzymes from other sources may be used; and other RNA polymerases with different promoter  
20 specificities may also be suitable for use.

Unless otherwise specified the reaction conditions used in the following examples were 40 mM Tris-HCl, 25 mM NaCl, 8 mM MgCl<sub>2</sub>, 5 mM dithiothreitol, 2 mM spermidine trihydrochloride, 1 mM rATP, 1 mM rCTP, 1 mM rGTP, 1 mM  
25 rUTP, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 0.15  $\mu$ M each primer or splice template, and specified amounts of template and enzymes in 100  $\mu$ l volumes.

These reaction conditions are not necessarily optimized, and have been changed as noted for some  
30 systems. The oligonucleotide sequences used are exemplary and are not meant to be limiting as other sequences have been employed for these and other target sequences.

Example 1Preliminary Procedure I

To demonstrate that this system worked, each of the four promoter containing oligonucleotides described above were each respectively put into reactions with or without 4 fmol target (frag 1). The reaction was incubated 2 minutes at 95°C to denature the target, then cooled to allow the oligonucleotides to anneal to the target. Reverse transcriptase, 13 Units, and T7 RNA polymerase, 100 Units, were added and the reaction was incubated 30 minutes at 37°C. One-tenth of the reaction was assayed using the hybridization method. The results (in Relative Light Units or "RLU(s)" and fmols) presented in Table 1 show that both the blocked and unblocked oligonucleotides serve as splice templates to produce product. The signals measured for reactions without target represent typical background levels of signal, for this type of assay.

Table 1. Comparison of Splice Templates in Preliminary Procedure I.

Spice Template	Target	Probe(-)	<u>Product</u>	
			RLU	fmol
ddT7pro(+)	4 fmol	Probe(-)	38451	240
ddT7pro(+)	0 fmol	Probe(-)	544	0
T7pro(+)	4 fmol	Probe(-)	65111	410
T7pro(+)	0 fmol	Probe(-)	517	0
ddT7pro(-)	4 fmol	Probe(+)	47756	85
ddT7pro(-)	0 fmol	Probe(+)	879	0
T7pro(-)	4 fmol	Probe(+)	156794	290
T7pro(-)	0 fmol	Probe(+)	600	0

40 Example 2Cycling with Preliminary Procedure I

The amplification system was cycled with ddT7pro(+) and T7pro(+). In this experiment, 4 amol frag I, HBV(-)Pr

and ddT7pro(+) or T7pro(+) were mixed in standard reactions and incubated at 95°C. After cooling, 13 Units of reverse transcriptase and 100 units of T7 RNA polymerase were added and the mixture was incubated 30 minutes at 37°C. One-tenth of the reaction was removed for assay and the cycle was repeated with the remainder. After repeating the cycle a third time, the 10 $\mu$ l aliquots were assayed by the hybridization method using Probe(-). The results presented in Table 2 indicate that product is amplified through cycling with both blocked and unblocked splice templates.

Table 2. Cycling with Preliminary Procedure I.

Splice template	Target	Relative Light Units(RLU's)		
		Cycle 1	Cycle 2	Cycle 3
ddT7pro(+)	4 amol	602	1986	10150
ddT7pro(+)	0 amol	658	592	595
T7pro(+)	4 amol	891	6180	52160
T7pro(+)	0 amol	496	504	776

25

### Example 3

#### Sensitivity of Preliminary Procedure I

In this example the unblocked splice template, T7pro(+), and the primer, HBV(-)Pr, were used to test the sensitivity of the amplification method. Six cycles of Preliminary Procedure I were run as described in Example 2 with decreasing quantities of frag 1. After amplification, the product was assayed using the hybridization method described in the Detailed Description of the Invention. Using this method, 4x10<sup>-21</sup> moles frag 1 could be detected (see Table 3).

Table 3. Sensitivity using 6 cycles of Preliminary Procedure I.

	Target (moles)	Sample $\mu$ l	Product RLU's
5	$4 \times 10^{-18}$	5	328390
	$4 \times 10^{-19}$	20	105364
	$4 \times 10^{-20}$	20	3166
	$4 \times 10^{-21}$	20	1927
	0	20	806

10

#### Example 4

##### Amplification Including Preliminary Procedure I

In the following example, the target to be amplified was frag 1. In the first set of reactions, various combinations of target and splice templates were incubated at 95°C for two minutes then cooled prior to adding 13 Units of reverse transcriptase and 100 Units of T7 RNA polymerase. The reactions were incubated 30 minutes at 37°C then 5 $\mu$ l aliquots of the reactions were assayed with both probes to quantitate the products. Subsequent to this assay, reactions were prepared using 5 $\mu$ l of reactions 1 and 2 and the T7pro(-) splice template. The mixtures were incubated 2 minutes at 95°C then cooled prior to adding 13 Units of reverse and 100 Units of T7 RNA polymerase. The new reactions were then mixed and incubated at 37°C for 2 hours. Aliquots of 10 $\mu$ l were removed to an ice bath at time points indicated in Table 4 below. The products were assayed using the hybridization method previously described. The data indicate that both splice templates allow production of RNA from frag 1. The data also indicate significantly more minus and plus sense products are produced in the reactions containing RNA and the splice template complementary to that RNA. And, finally, the reaction kinetics for the reaction 1B show a geometric increase in product whereas the kinetics for the 2B reaction are of a more linear form. This difference

indicates the product in the 1B reaction is serving as a substrate to generate more product; thus the reaction is autocatalytic.

5 Table 4. Preliminary Procedure I

	Reaction		1	2	3	4
10	Target (4 fmol)		Yes	Yes	Yes	No
	T7pro(+)		Yes	No	No	No
	T7pro(-)		No	No	Yes	No
	Probe	Time	Relative Light Units(RLU's)			
15	Probe(+)	30'	1156	1058	21859	591
	Probe(-)	30'	11693	771	744	691
	Reaction		1A	1B	2A	2B
20	Reaction1		Yes	Yes	No	No
	Reaction2		No	No	Yes	Yes
	T7pro(-)		No	Yes	No	Yes
	Probe	Time	Relative Light Units			
25	Probe(+)	0'	714	757	639	661
		30'	686	339	1663	1373
		60'	718	6331	645	1786
		120'	816	16889	660	2238
30	Probe(-)	120'	3142	6886	637	780

### 35 Example 5

#### Reaction Kinetics for Amplification Including Preliminary Procedure I

The following example further demonstrates the potential of this embodiment of the methods the invention. A small quantity of frag 1 (10 amol), was used in reactions with various combinations of T7pro(+), T7pro(-), and HBV(-)Pr. The reaction mixtures were incubated at 95°C for two minutes to denature the DNA target and cooled prior to adding reverse transcriptase and T7 RNA polymerase. After mixing, the reactions were incubated 30 minutes at 37°C. Fifteen microliter aliquots were removed at various time points and stored at 0°C until assayed.

The hybridization assay was used to quantitate the products of the reactions. The data presented in Table 5 show that the invention requires one splice template and one primer. A second splice template is advantageous, however. The results with only one primer or splice template were below the detection limits of the assay. As in the previous example, the reaction kinetics are geometric, indicating an autocatalytic system.

Table 5. Preliminary Procedure II Reaction Kinetics.

Reaction	1	2	3	4	5	6	7
Target (10 amol)	No	Yes	Yes	Yes	Yes	Yes	Yes
T7pro(+)	Yes	No	Yes	No	No	Yes	Yes
T7pro(-)	Yes	No	No	Yes	No	No	Yes
HBV(-)Pr	Yes	No	No	No	Yes	Yes	No
Time (minutes)	Minus Product (RLU's)						
0	619	638	635	703	592	619	656
30	613	635	613	755	626	844	1133
60	635	649	856	894	635	2146	6008
90	593	619	619	925	624	6226	23484
120	621	606	627	946	639	12573	43939
180	678	635	714	930	627	21719	78682
Time (minutes)	Plus Product (RLU's)						
0	624	646	1661	710	621	636	962
30	637	601	802	629	655	803	758
60	639	706	800	679	664	226	2895
90	638	683	956	633	687	7786	8085
120	643	670	884	647	632	18160	18241
180	683	617	968	758	712	34412	41165

Our subsequent work has demonstrated continued product synthesis for over 5.0 hours, which is substantially better than prior art methods. We also have demonstrated increased sensitivity.

Example 6Amplification Including Preliminary Procedure II

In this example various combinations of primers were used to amplify 500 amol of a DNA target without defined termini. The target was the M13L(-) referenced above. Upon reaction preparation, the samples were incubated two minutes at 95°C, then cooled prior to adding 13 Units of reverse transcriptase. The reactions were then incubated twelve minutes at 42°C. Next the reactions were again heated for two minutes at 95°C and cooled. Reverse transcriptase and T7 RNA polymerase were added and the reactions were incubated for two hours at 37°C. Ten microliter aliquots of the reaction were assayed with both Probe(+) and Probe(-). The results presented in Table 6 show that synthesis of large amounts of nucleic acid occurs only when two primers are employed. They also demonstrate the benefit of two promoter-primers over one promoter-primer and one primer. The low level synthesis in reactions 4 and 5 correspond to synthesis of approximately one copy of DNA from the original template. The system employs an initial 95°C denaturation step which may serve to denature double-stranded targets or double-stranded regions of a single-stranded target as well as inactivate unwanted nuclease and protease activities.

Table 6. Preliminary Procedure II System.

Reaction	1	2	3	4	5	6	7
M13L(-)	No	No	Yes	Yes	Yes	Yes	Yes
T7pro(+)	Yes	Yes	No	Yes	No	Yes	Yes
T7pro(-)	Yes	No	No	No	Yes	Yes	No
HBV(-)Pr	No	Yes	No	No	No	No	Yes
Probe	Relative Light Units (RLU's)						
Probe(+)	862	744	762	1089	2577	96221	30501
Probe(-)	473	420	483	3038	1080	15171	14863

Example 7Effect of RNase H

To demonstrate that the addition of exogenous RNase H may improve amplification in the autocatalytic systems described of the present invention, several reactions were prepared using various quantities of target, M13L(-), and either 0 or 2 Units of exogenous RNase H. The exogenous RNase H used was derived from *E. coli*. All reactions were prepared with T7pro(+) and T7pro(-). The reactions were subjected to the 95°C denaturation and cooled prior to adding reverse transcriptase. After a twelve minute incubation at 42°C, the reactions were again denatured at 95°C. After cooling, reverse transcriptase, T7 RNA polymerase, and, if indicated (see Table 7), RNase H was added and the reactions were incubated for 3 hours at 37°C. Aliquots of 10μl were removed from each reaction at hourly intervals for assay by the hybridization method. The data in Table 7 show that exogenous RNase H significantly enhanced the reaction kinetics and increases the sensitivity of the invention. Signals less than 600 RLU's were interpreted as typical background levels.

Table 7. Effect of Exogenous RNase H on Amplification Including Preliminary Procedure II Sensitivity.

	Target (moles)	RNase H (Units)	RLU's at Time(hours)			
			0	1	2	3
30	1x10 <sup>-17</sup>	0	478	7659	28716	60443
	1x10 <sup>-18</sup>	0	440	946	13332	36039
	1x10 <sup>-19</sup>	0	413	581	10063	41415
	1x10 <sup>-20</sup>	0	406	424	717	4520
35	0	0	455	376	579	2075
40	1x10 <sup>-18</sup>	2	419	20711	50389	64073
	1x10 <sup>-19</sup>	2	411	6831	21312	29818
	1x10 <sup>-20</sup>	2	420	604	1281	1375

Example 8Effect of Exogenous DNA

It has been demonstrated that the addition of exogenous DNA may significantly inhibit this autocatalytic amplification system. To further demonstrate the benefit of adding exogenous RNase H, amplification reactions were prepared with or without 2  $\mu$ g calf thymus DNA to demonstrate this inhibition. In reactions with the calf thymus DNA, two concentrations of reverse transcriptase were employed to test whether additional AMV RNase H would overcome the inhibition. Also, RNase H from E. coli was added to some of the reactions for the same reasons. The reactions differ from the standard reactions in that the concentration for each of the ribonucleotides were increased to 2.5 mM and the concentration of magnesium chloride was increased to 12.8 mM. The reactions were prepared using 100 amol of M13L(-) as a target and T7pro(+) and T7pro(-). After denaturing two minutes at 95°C, the reactions were cooled and 13 or 39 Units of reverse transcriptase were added and the reactions were incubated 12 minutes at 37°C. The reactions were again denatured at 95°C and cooled prior to adding 13 or 39 Units of reverse transcriptase, 100 or 300 units of T7 RNA polymerase, and either 0 or 2 Units of E. coli RNase H. After incubating one hour at 37°C, 10  $\mu$ l of each reaction was assayed using the hybridization assay. The results presented in Table 8 showed that the calf thymus DNA inhibited the reaction by 90% in comparison to a reaction system without exogenous DNA and that additional reverse transcriptase (and its associated RNase H) did not significantly affect the product amplification. The addition of more T7 RNA polymerase did have a significant effect on the product yield, but it was small relative to the increase due to addition of exogenous RNase H. Not only was the inhibition eliminated, the amplification was increased over five-fold relative to the reaction without the calf thymus DNA and E. coli RNase H. The signals

observed with the higher amount of E. coli RNase H were saturating for the amount of probe used in the hybridization assay. To accurately quantitate these samples, dilution of the amplified product would be required before  
5 assaying.

Table 8. Effect of RNase H on Amplification  
Inhibition by Exogenous DNA

	Reverse	T7 RNA	E. coli	Target	Exogenous	Relative
	Transcriptase	Polymerase	RNase H	DNA	DNA	Light
	(Units)	(Units)	(Units)	(amol)	( $\mu$ g)	Units
10	13	100	0	0	0	458
	13	100	0	100	0	55305
	13	100	0	100	2	3003
15	39	100	0	100	2	2786
	13	300	0	100	2	5434
	39	300	0	100	2	6831
	13	100	4	100	2	278666
	39	100	4	100	2	334649
20	13	300	4	100	2	359101
	39	300	4	100	2	375043

#### Example 9

#### 25 Amplification By the General Method

This system does not require an initial transcription and denaturation; a DNA complement of the target is made and the original target is removed by the RNase H. The DNA may then anneal to a second primer or promoter-primer  
30 and through DNA synthesis produce a template for the RNA polymerase. If the RNase H is not active, the DNA:RNA hybrid produced first will end the reaction without producing a template for the RNA polymerase. In an attempt to demonstrate the method of this invention, a  
35 plasmid containing an RNA polymerase promoter and the target nucleic acid was used to produce large quantities of single-stranded RNA transcripts containing the target

sequence within other sequences. Two similar reactions were also prepared: one containing the plasmid without the RNA polymerase and the other with the RNA polymerase without the plasmid.. Dilutions of each of these reactions were assayed to quantitate the products. Equivalent dilutions of all three reactions were used to prepare amplification reactions containing the two promoter-primers, T7 pro(+) and T7 pro(-). Reaction 1 of Table 9 contained 60 amol of RNA and 0.6 amol of the starting plasmid. Reaction 2 contained 0.6 amol of the starting plasmid, but no RNA. Reaction 3 contained no target. The reactions were not denatured at 95°C; instead reverse transcriptase and T7 RNA polymerase were added and the reaction was incubated at 37°C for four hours. Aliquots were removed hourly for later assay by the hybridization method. As shown in Table 9, the reaction containing the plasmid and the RNA produced from the plasmid gave large hybridization signals; the reaction containing the plasmid alone produced significant product as the T7 RNA polymerase could produce RNA from the plasmid which could then be utilized according to the General Method to produce more nucleic acid of both senses; and, finally, the control (Reaction 3) containing no target produced nothing.

Table 9. Preliminary Procedure IV Reaction Kinetics.

Time (hours)	Reaction 1 RLU's for Probe(+)	2 1890	3 886	1 7979	2 1074	3 434
2	9695	1890	886	7979	1074	434
3	22266	2819	824	15018	1487	491
4	33863	4571	828	16540	2310	556

Rxn 1: plasmid plus RNA.

Rxn 2: plasmid.

Rxn 3: no target

Example 10Amplification by the General Method

The following experiment was done to determine if other methods of initiation were possible for the invention which may alleviate the need for the first primer extension and denaturation step for DNA targets without defined termini. In this experiment, the stated quantities of target are amplified with or without the first reverse transcriptase step as indicated in Table 10.

10 The amplification time once all enzymes are added was four hours. The target used was the M13(+) diluted into normal saline. T7pro(+) and T7pro(-) were used. To 25  $\mu$ l of each dilution was added 25  $\mu$ l 0.1N KOH. The samples were incubated ten minutes at 98°C and then cooled. Each

15 sample was brought to 100  $\mu$ l and final concentrations of 50 mM Trizma base, 40 mM glutamic acid, 25 mM potassium hydroxide, 12.8 mM magnesium chloride, 5 mM dithiothreitol, 2 mM spermidine trihydrochloride, 2.5 mM each ribonucleotide triphosphate, 0.2 mM each deoxyribo-

20 nucleotide triphosphate, and 0.15  $\mu$ M each primer. To the Standard Protocol tubes, 13U reverse transcriptase was added and the reactions were incubated 12 minutes at 42°C, then two minutes at 95°C, and cooled. Then to all tubes was added 13U reverse transcriptase, 100U T7 RNA

25 polymerase, and 0.5U RNase H. The reactions were then incubated four hours at 37°C and 10  $\mu$ l aliquots were assayed using the chemiluminescent probe assay. The results presented in Table 10 show that some amplification was evident using the abbreviated protocol. And although

30 the level of amplification observed was significantly less than that for the Standard Protocol, this may be further developed to be as efficient or may be useful in cases where significant levels of target nucleic acid are present.

Table 10. Amplification Without First Primer Extension

	Target	Moles	Protocol	RLU's
	M13(+)	3.8E-17	Standard	2295451
	"	3.8E-19	"	2374443
5	"	3.8E-21	"	230227
	Negative	0	"	3037
	M13(+)	3.8E-17	Short	2475574
	"	3.8E-19	"	27209
10	"	3.8E-21	"	17144
	Negative	0	"	1679

The likely explanation for these data is that T7RNA polymerase is not completely specific. There is a low level of random RNA synthesis which generates small numbers of RNA copies of the target regions. These copies are amplified via the standard method.

\* \* \* \*

Our initial work demonstrated excellent amplification with certain primer sets and targets without the addition of exogenous RNase H. Our subsequent work clarifying the mechanism of the reaction has made it possible to efficiently apply the method to a wider variety of targets. We disclose and claim herein methods which both use exogenous RNase H in amplification and those which rely on RNase H activity associated with reverse transcriptase.

We have discovered that E. coli RNase should not routinely be added as it does not always improve amplification. We have determined that some forms of RNase H are sequence specific as to where they cut the RNA of RNA:DNA hybrids. In the amplification reaction, we have not detected the promoter-containing primer in full-length product DNA. In embodiments using two promoter-primers, only one of the promoter primers is detectably incorporated into full-length product DNA. All other mechanisms that have been postulated by those skilled in the art show full-length product DNA containing both

primers. We attribute these findings to the likelihood that RNase H is not fully degrading the major RNA species synthesized during amplification. Based on these findings, a new amplification mechanism is set forth herein which incorporates our findings regarding RNase H sequence specificity. New and useful promoter-primer design criteria are disclosed. Furthermore, we claim herein novel methods for synthesizing multiple copies of a target nucleic acid sequence, comprising

10       1) selecting a primer, complementary to a portion of the RNA target sequence, which complexes with the portion of the RNA target, said portion of the target located such that it remains capable of forming primer extension product after being exposed to degradation by a  
15       selected RNase H;

          2) selecting a promoter-primer complementary to a portion of the DNA to be obtained by extension of the primer, which complexes with the DNA in an area where substantially all of the complementary RNA is removed from  
20       the duplex through degradation of RNase H; and

          3) combining the RNA target with the primer, promoter-primer, RNase H, reverse transcriptase and transcripase and forming multiple copies of the RNA and multiple copies of DNA complementary to the RNA. The  
25       novel methods herein described do not make a substantially equivalent number of copies of DNA of the same polarity (or "sense") as the RNA target sequence.

          This procedure provides a method which permits the design of efficient promoter-primers and primers for new  
30       target sites. We disclose and claim herein such promoter-primer and primer combinations and design criteria for same. The mechanism disclosed herein involves a novel reaction intermediate for transcription of RNA strands. This intermediate is a double-stranded complex of RNA and  
35       DNA which contains a double-stranded DNA promoter region. Nothing like this has, to our knowledge, ever been described in the literature. Indeed, none of the prior

art systems, specifically neither Guatelli, J.C. et al.,  
87 PNAS 1874-1878 (1990), nor PCT App. Ser. No. 88/10315  
to Gingeras, T.R. et al., properly select the primer and  
promoter-primer sequence based upon the location of RNase  
5 H degradation sites. Thus, the methods herein are novel  
and nonobvious from any previously disclosed.

Recognition of the importance of the RNase H sequence  
specificity and an understanding of the reaction mechanism  
is key to the efficient application of this target  
10 amplification method to a wide variety of target  
sequences. Moreover, until this time, practitioners  
assumed that RNase H fully and systematically degraded the  
RNA strand of an RNA:DNA complex.

15 I. The Mechanism of Amplification Methods

Testing amplification efficiency with both prior art  
methods revealed a great deal of variability in  
amplification efficiency with small changes in the primer  
sets used. The reaction method generally accepted in the  
20 prior art did not, in our view, provide a reasonable  
explanation as to why one primer set worked so much better  
than another.

Attempts to improve amplification using the prior art  
methods did not give satisfactory results. Efficiency of  
25 priming was examined to see if differences in the ability  
to initiate DNA chains were responsible for observed  
differences in primer set efficiency. No correlation  
between priming efficiency and overall amplification could  
be found. Analysis of primer sets for the ability to form  
30 self-complementary structures and cross-complementary  
structures indicated that differences in primer efficiency  
were not solely attributable to these factors either.

We also found that the addition of E. coli RNase H  
did not uniformly improve amplification. As the data  
35 submitted herein show, the results observed varied from  
target to target and from primer set to primer set. The  
amount of E. coli RNase H added also is v ry important and

must be kept within a narrowly defined range. With a given target and primer set, addition of *E. coli* RNase H is helpful in some cases when the reverse transcriptase is that from avian myeloblastosis virus ("AMV") but not  
5 when the reverse transcriptase is that from Moloney murine leukemia virus ("MMLV"). Data illustrating these conclusions are provided herein.

Earlier work suggested that AMV reverse transcriptase leaves relatively large fragments when it digests the RNA  
10 from the RNA:DNA hybrid formed during virus replication. These fragments serve as primers to initiate synthesis of the second DNA strand. We report herein our findings that there is evidence for sequence specificity of the AMV and MMLV RNase H activities.

15 In order to elucidate the mechanism of the reaction, individual primers were terminally labeled with  $^{32}\text{P}$ , and the incorporation of each primer into DNA products was examined by polyacrylamide gel electrophoresis. According to the generally accepted prior art mechanism, both  
20 primers would be incorporated into full length DNA products. Our experiments showed, however, that only the primer complementary to the major RNA species synthesized during amplification was incorporated into full length product. The primer having the same polarity as the major  
25 RNA strand was never detected in full length DNA product. In fact, it remained quite small. These results were confirmed with a number of different targets and primer sets.

The failure to detect extension of one of the primers  
30 indicated that a fully double-stranded DNA intermediate did not accumulate during amplification, and was not required for autocatalytic amplification. These observations indicate a mechanism for the amplification systems of this invention which takes into account  
35 probable sequence specificities of the RNase H. The mechanism is generally depicted in Fig. 4.

Experiments have shown that the enzyme cuts the RNA of an RNA:DNA hybrid into specific pieces. Furthermore, the locations of the cut sites were shown to be in specific regions. To confirm the mechanism, an RNA:DNA  
5 hybrid was prepared which contained the plus strand RNA that would be generated from our T7pro<sup>+</sup>/T7pro<sup>-</sup> target and primer set combination. The <sup>32</sup>P labelled RNA was hybridized to complementary DNA, incubated with AMV reverse transcriptase (containing its associated RNase H  
10 activity), and the reaction products were analyzed by polyacrylamide gel electrophoresis (Figure 5). The fragment size indicated that several small pieces were generated and that these were produced by cuts near one or both ends. The interior region of the molecule was not  
15 cut. This experiment demonstrates that the enzyme has sequence or structural specificity under the reaction conditions used. The results were entirely consistent with the reaction mechanism of Fig. 4.

Further experiments were performed to determine where  
20 the cut sites occurred. It is preferred that multiple cut sites occur in the region homologous to the promoter-containing primer and not in the region binding the other primer. By labeling the termini of the RNA individually and analyzing the digestion products, it was found that  
25 under the conditions used the cuts were detected only at the 5' end of the RNA. This is consistent with the mechanism of Fig. 4.

Sequencing experiments were performed to determine the sequences at which the RNase H activities of AMV and  
30 MMLV reverse transcriptases cut. Sequences were identified that were specifically cut by each enzyme. As predicted, the sequence specificities of the two enzymes are different, and this is believed to explain why some primer sets work better with one enzyme and some with  
35 another. The sequence specificities obtained for the MMLV enzyme under our reaction conditions do not match those reported in the literature under another set of reaction

conditions, indicating that specificity may be influenced by reaction conditions.

Scrutiny of the role of the RNase H in the amplification mechanism has resulted in our finding that completely removing the promoter directed transcript from its cDNA copy may not be necessary, or even desirable for formation of a new transcriptionally active template. Thus, in some applications, even a very low level of RNase H activity, deriving from the reverse transcriptase-intrinsic RNase H, will be sufficient for effective amplification if the RNase H is more site selective in allowing the promoter-primer to anneal to the first strand cDNA product or if it interferes less with the annealing of the other primer to the transcript.

Since E. coli RNase H is reportedly less specific than the retroviral enzymes, it may cleave in the region to which the non-promoter containing primer binds, especially if the concentrations of this primer, the target, the E. coli RNase H, and components affecting the enzyme activity are not carefully balanced. In our view these results make the use of E-coli RNase H non-preferable in commercial applications. Addition of another RNase H activity, one with different specificities, may be useful in those cases in which the reverse transcriptase RNase H does not cut in the desired regions or does not cut under appropriate conditions. Work with MMLV reverse transcriptase, for example, has shown that this enzyme is less sensitive than the AMV enzyme to inhibition by sample DNA. It is the best mode for many systems.

New primer sets were designed and are set forth herein based upon the model and the RNase H sequence specificity information that we have obtained to date. Significantly better synthesis was obtained from these primer sets than was obtained with those designed previously without knowledge of the mechanism and sequence specificities. The invention herein described makes

possible the design of functional primer sets for specific target regions.

The new mechanism we have discovered involves a novel reaction intermediate for transcription of RNA strands.

5 This intermediate is a double-stranded complex of RNA and DNA which also contains a double stranded DNA promoter region. To our knowledge, the reaction is demonstrably different from any previously disclosed.

An understanding of the reaction mechanism is  
10 critical to using these target amplification procedures. Recognition of the importance of the RNase H sequence specificity is key to the efficient application of this target amplification method to a wide variety of target sequences. On the other hand, the empirical approach to  
15 promoter-primer design is very intensive, costly, and has a low frequency of success, making this invention a useful advance in the art.

A. Narrow Range of Activity for RNase H Concentration

20 The amplification system with E. coli RNase H initially was considered to be the preferred embodiment because greater synthesis was achieved with the particular target and primer set being studied, and the E. coli RNase H was found to be useful in helping to overcome inhibition  
25 by sample DNA. However, analysis of the reaction indicates that addition of E. coli RNase H is detrimental to amplification in many cases. Moreover, amounts of E. coli RNase H added must be carefully controlled over a narrow range since the presence of too much or too little  
30 is harmful. For practical commercial application of the method, this is a significant drawback, especially since the enzyme may not be completely stable on storage. In addition, the use of E. coli RNase H adds significant cost and complexity to the system. The cost may be prohibitive  
35 for many commercial applications. Using E. coli RNase H makes the assay more complex, which in turn increases research and development costs and may make the assay too

delicate for wide commercial applicati n. Thus, our elucidation of the assay mechanism has resulted in m thods for a widely applicable assay both in terms of technical feasibility (applicability to target sites) as well as  
5 being a cheaper and a more robust procedure. Since the addition of E. coli RNase H was found to result in increased amplification in early experiments, the effect of E. coli RNase H on the performance of the amplification system in samples containing serum or human DNA was  
10 examined in several experiments.

#### Example 11

##### Optimization of E. coli RNase H Concentration

Experiments were performed to determine the amount of  
15 E. coli RNaseH needed for optimal amplification in serum. The following experiment compared amplification with the T7pro<sup>+</sup>/T7pro<sup>-</sup> primer pair in the presence of 0, 0.25, 0.5 and 1 U of RNaseH per assay. HBV + plasma diluted to the levels shown in HBV- human serum or HBV- serum alone was  
20 tested.

Ten  $\mu$ l of serum were added to an equal volume of 0.1 N KOH and covered with a layer of oil to prevent evaporation. The samples were mixed, heated at 95°C and allowed to cool to room temperature. The samples were  
25 brought to 90  $\mu$ l reaction volume with a final concentration of 50 mM Tris acetate pH 7.6, 20.8 mM MgCl<sub>2</sub>, 5 mM dithiothreitol, 2 mM spermidine hydrochloride, 0.15  $\mu$ M each primer, 6.25 mM GTP, 6.25 mM ATP, 2.5 mM UTP, 2.5 mM CTP, 0.2 mM each dTTP, dATP, dGTP, dCTP and 13 U of AMV  
30 reverse transcriptase. The samples were mixed and heated at 37°C for 12 minutes, then heated to 95°C and cooled to room temperature. Thirteen units of RT and 100 U of T7 RNA polymerase were added and the reactions were incubated for three hours at 37°C. Twenty-five  $\mu$ l of each reaction  
35 was assayed.

The data show that ther is a narrow optimum range of concentration of E. coli RNase H centering around 0.25 U

E. coli RNase H per reaction for this system. Even though E. coli RNase H is difficult to use, some added RNase H activity was beneficial in this experiment.

5		Table 11	
	Moles Target	RNaseH (Units)	RLU observed
	$5 \times 10^{-20}$	0	567809
	$5 \times 10^{-22}$		18041
10	$5 \times 10^{-23}$		2938
	0		1634
	$5 \times 10^{-20}$	0.25	1153366
	$5 \times 10^{-22}$		732109
15	$5 \times 10^{-23}$		5566
	0		1423
	$5 \times 10^{-20}$	0.5	1001904
	$5 \times 10^{-22}$		29596
20	$5 \times 10^{-23}$		1793
	0		1898
	$5 \times 10^{-20}$	1.0	610485
	$5 \times 10^{-22}$		13026
	$5 \times 10^{-23}$		4062
25	0		1662

#### Example 12

Next the amount of E. coli RNase H needed for optimal amplification of an HIV primer pair was determined in the presence or absence of a lysate containing 8  $\mu$ g of human DNA.  $2 \times 10^{-18}$  moles of viral target were present in each reaction. DNA target was mixed with 50 pmol of each primer in 40 mM Tris HCl pH 8.3, 25 mM NaCl, 20.8 mM MgCl<sub>2</sub>, 5 mM dithiothreitol, 2 mM spermidine hydrochloride, and nucleotide triphosphates as described for Table 11, heated to 95°C and cooled to room temperature. Thirteen units of AMV reverse transcriptase were added and the reaction

heated to 42°C for 12 minutes, to 95°C and cooled again to room temperature. Thirt en units of AMV reverse transcriptase and 100 units of T7 RNA polymerase were added and the reactions heated to 37°C for 3.5 hours prior  
5 to assay.

Table 12

	<u>Lysate</u>	<u>RNase H</u>	<u>RLU</u>
10	-	0 U	8,400
	-	0.5	239,000
	-	1.0	468,000
	-	1.5	498,000
	-	2.0	439,000
15	-	3.0	20,100
	-	4.0	5,806
	+	.0	1,667
	+	0.5	924
	+	1.0	6,635
20	+	1.5	579
	+	2.0	13,400
	+	3.0	17,800
	+	4.0	9,152

25 These results illustrate that E. coli RNase H levels have to be carefully controlled as too much E. coli RNase H was detrimental to the amplification. Additionally, the optimal concentration was altered by the presence of non-specific human DNA and the inhibition by human DNA was  
30 significant at all RNase H levels.

### Example 13

We investigated the effect of E. coli RNase H on amplification of a second region referred to as HIV region  
35 2. The following data demonstrate that E. coli RNase H enhances amplification within a narrow concentration range. The HIV region 2 primers were amplified as

described for Table 12 in the presence of different concentrations of E. coli RNase H. Ten microliters of each reaction were assayed and dilutions were made when necessary. Signals were compared to a standard curve to determine the amount of target made.

	RNase H	pmole target	pmole product	Amplification observed
	-	0	0	0
	-	$1.67 \times 10^{-10}$	2.14	$1.3 \times 10^{10}$
10	0.2 U	$1.67 \times 10^{-10}$	0.17	$1.0 \times 10^9$
	0.4 U	$1.67 \times 10^{-10}$	0.18	$1.1 \times 10^9$
	1.2 U	$1.67 \times 10^{-10}$	0.012	$7.6 \times 10^7$
	-	$1.67 \times 10^{-8}$	16.0	$9.6 \times 10^8$
15	0.2 U	$1.67 \times 10^{-8}$	20.6	$1.2 \times 10^9$
	0.4 U	$1.67 \times 10^{-8}$	0.14	$8.5 \times 10^6$
	1.2 U	$1.67 \times 10^{-8}$	0.15	$9.0 \times 10^6$

These data show that amplification can be achieved without the addition of E. coli RNase H, contrary to the assertions of Guatelli, et al., 87 PNAS 1874-1878.

We investigated using E. coli RNase H with MMLV reverse transcriptase in several target regions. Reactions were done in the presence or absence of 8  $\mu$ g human DNA using conditions described for Example 12, with a 3 hour autocatalysis step. Primer sets from HIV regions 1, 3 and 4 were tested. The amount of viral template used was selected to give RLU in the linear range of the hybridization assay. MMLV reverse transcriptase was used at 400 U during initiation, 800 U for efficient amplification. 400 U of T7 RNA polymerase were included, and 1 U of E. coli RNase H was added as indicated. Values presented under the column headings, +RNase H, -RNase H, are RLUs obtained from assay of 10  $\mu$ l of the reactions.

Table 15

Target		Human		
Region	Moles Target	DNA	+RNase H	-RNase H
HIV region 1	$2 \times 10^{-21}$	-	54,900	137,000
5 HIV region 1	$2 \times 10^{-21}$	8 $\mu$ g	15,100	13,800
HIV region 3	$2 \times 10^{-20}$	-	96,100	391,000
HIV region 3	$2 \times 10^{-20}$	8 $\mu$ g	124,000	246,000
HIV region 4	$2 \times 10^{-21}$	-	20,400	107,000
HIV region 4	$2 \times 10^{-21}$	8 $\mu$ g	56,000	8,800

10

In the presence of DNA, E. coli RNase H apparently stimulated amplification directed by the HIV region 4 primers. In most cases we have tested, amplification using MMLV reverse transcriptase alone is at least as good as when MMLV reverse transcriptase is used with E. coli RNase H. E. coli RNase H is not required for efficient amplification, contrary to the assertions of Guatelli, et al., 87 PNAS 1874-1878.

#### 20 Sequence Specificity of Reverse Transcriptase

We also have discovered that some primer sets work best with AMV reverse transcriptase while others work best with MMLV reverse transcriptase or with one of the reverse transcriptases and added E. coli RNase H. The observed degree of variability in amplification efficiency with small changes in promoter primers and primers or source of RNase H supports our proposed mechanism. We set forth below our detailed data in support of these findings.

MMLV reverse transcriptase has been cloned, and is commercially available from BRL (Bethesda Research Labs), U.S. Biochemicals and others in a very concentrated form (greater than 300 units per  $\mu$ l). It should be noted that comparable DNA synthetic activity on natural nucleic acid templates is obtained with approximately 10-fold greater unit concentration of MMLV reverse transcriptase compared

35

to AMV reverse transcriptase. Lack of comparability in unit activity is due to the fact that the enzymes show different relative activities when tested with homopolymer templates (used in the unit activity assay) and heteropolymeric nucleic acid templates. We tested the use of MMLV reverse transcriptase at various levels in our amplification reactions. Examples of these results are shown in the tables below. In the AMV reverse transcriptase samples, reverse transcriptase was used at 14 U during the initiation step and 56 U during the amplification step. The amount of MMLV reverse transcriptase was titrated for both the initiation and the amplification steps. The incubation conditions used were as described for Example 12 except that 15 pmol of each HIV region 2 primer was used and 25 mM KCl replaced the NaCl. T7 polymerase was used at 400 U during the amplification. The following table shows performance in the presence or absence of 8  $\mu$ g human DNA. Columns headed with the designation AMV or MMLV show the results of amplifications performed with AMV reverse transcriptase or MMLV reverse transcriptase, respectively. The numbers refer to the number of units used during initiation and autocatalysis, respectively. The values contained within the table are RLUs. Note that dilutions of the amplification products were not performed and values >200,000 RLU may significantly underestimate the extent of amplification since signal saturation occurs at a level of hybridization target sufficient to give about 250,000 RLU with the conditions used.

Table 16

		Human	AMV	MMLV	MMLV	MMLV
	Moles Target	DNA	14/56	400/400	400/600	400/800
	0	-	495	470	-	3,800
5	$1.6 \times 10^{-22}$	-	278,000	77,000	-	5,621
	$1.6 \times 10^{-20}$	-	292,000	278,000	-	269,000
	0	+	474	547	488	1,352
	$1.6 \times 10^{-20}$	+	10,200	62,700	205,000	149,000

10 Although the sensitivity of amplification directed by MMLV reverse transcriptase in the absence of human DNA was significantly lower than AMV directed amplification, the MMLV was much more effective in the presence of exogenous DNA.

15 After observing the high level amplification of the HIV region 2, we tested the other target regions in the presence of human DNA and found that, using AMV reverse transcriptase, E. coli RNase H was still required for the most effective amplification in these regions. We then  
 20 tested each target region using MMLV reverse transcriptase, without E. coli RNase H, to compare amplification performance with reactions containing AMV reverse transcriptase + E. coli RNase H. An example of these results for two target regions is shown in the table  
 25 below. The HIV region 3 and 4 primers were used (50 pmol per reaction) as described for Example 12. In reactions using AMV reverse transcriptase, 14 U was used at initiation, 56 U reverse transcriptase + 1 U E. coli RNase H were added for amplification. In reactions using MMLV  
 30 reverse transcriptase, 400 U was added at initiation and 800 U for amplification. All reactions contained 400 U T7 RNA polymerase during the four-hour amplification incubation. Values within the tables are RLU obtained from Homogeneous Protection Assay performed using 10  $\mu$ l of the  
 35 amplification reactions. Dilutions of the reactions were performed in some cases before assay.

68

Table 17

	Moles Target	Human DNA	HIV region 3		HIV region 4	
			AMV	MMLV	AMV	MMLV
5	0	-	2,049	751	1,184	777
	$1.6 \times 10^{-21}$	-	70,800	689	$2.1 \times 10^7$	305,000
	$1.6 \times 10^{-20}$	-	510,000	1,869	-	-
10	0	+	551	400	1,058	1,182
	$1.6 \times 10^{-21}$	+	-	-	13,900	16,200
	$1.6 \times 10^{-20}$	+	706	1,862	141,000	154,000
	$1.6 \times 10^{-19}$	+	-	-	683,000	723,000
	$1.6 \times 10^{-18}$	+	10,800	115,000	-	-

- 15 As observed in the HIV region 2, in the absence of human DNA, the amplification with MMLV is significantly less than with AMV reverse transcriptase with E. coli RNase H but in the presence of DNA the MMLV directed amplification is at least as good as accomplished by AMV reverse
- 20 transcriptase with RNase H.

#### Example 14

#### Primers for Second Region of HBV genome

- The following experiment was performed with primer
- 25 sets directed to two regions of the HBV genome. The data show that the primer sets do not amplify to the same extent with AMV RT. The experiment was performed as described for Example 11 using HBV positive plasma diluted in negative serum. Ten microliters of amplification
- 30 reaction were tested in the hybridization assay.

69

Table 18

RLU observed

	Moles Target	HBV Region 1	HBV Region 2
	$4.8 \times 10^{-21}$		690,674
5	$4.8 \times 10^{-22}$	475,849	73,114
	$4.8 \times 10^{-23}$	242,452	4,193
	0	1,417	1,940

These results were confirmed by additional experiments  
 10 using standard protocols. The region 1 primers  
 consistently gave higher RLU in these experiments.

In contrast, when 800 U of MMLV enzyme were used to  
 amplify the same two primer pairs, the opposite effect was  
 seen as shown below.

15

Table 19

RLU observed

	Moles Target	Region 1	Region 2
	$9.6 \times 10^{-20}$	37,278	1,067,951
20	$9.6 \times 10^{-22}$	1,858	40,826
	0	1,010	1,646

In this experiment, each reaction contained 5  $\mu$ l serum.

Thus, the amplification potential of each primer pair  
 25 was influenced by the reverse transcriptase present during  
 amplification. Factors such as the availability of the  
 template sequences, ability of the primers to hybridize  
 specifically, and efficiency of RNA synthesis should not  
 be affected significantly by the type of reverse  
 30 transcriptase present. Factors such as RNase H  
 specificity and activity and DNA polymerizing activity  
 could be affected.

The following data illustrates that promoter-primer combinations used in the appropriate conditions can be designed to increase amplification.

5

Table 21

This experiment was performed with HBV region 2 primers as described for Table 11 except that the entire amplification reaction was analyzed by hybridization.

	Target Molecules	Moles Target	RLU Observed
10	1200	$2 \times 10^{-21}$	1,094,177
	120	$2 \times 10^{-22}$	442,137
	12	$2 \times 10^{-23}$	24,053
	1.2	$2 \times 10^{-24}$	8,654
	0	0	1,828

15

Example 15Comparison of AMV reverse transcriptase and MMLV reverse transcriptase.

The following experiment compared amplification of primers for a BCL-2 chromosomal translocation major human chromosomal breakpoint t(14;18) found in patients with CML using MMLV (300 units) or AMV (39 units). The effect of E. coli RNase H was evaluated with each enzyme. Amplifications were performed as described for Example 12 except that 24 mM MgCl<sub>2</sub> and 50 pmol each primer were used. In reactions containing lysate, 4 µg of DNA from white blood cells was present. All reactions contained 300 units T7 RNA polymerase and 10 amol of input double-stranded DNA target.

30

Table 22

	RT	Lysate	0 Units RNase H	0.5 Units RNase H	1.0 Units RNase H	2.0 Units RNase H
	AMV	-	108,018	2,035,377	-	-
35		+	44,485	204,894	165,972	136,647
	MMLV	-	3,015,948	2,224,666	-	-
		+	3,070,136	2,714,257	767,218	105,845

The results show that MMLV and AMV RT do not amplify this primer set to the same extent, particularly in the absence of E. coli RNase H. E. coli RNase H added to reactions  
5 containing AMV reverse transcriptase markedly improved amplification; this indicates that the RNase activity was limiting in these reactions. In contrast, E. coli RNase H did not enhance amplification when added to reactions containing MMLV RT. The data also confirms a point  
10 already made concerning the ability of MMLV RT to sustain significant amplification in the presence of large amounts of nonspecific human DNA.

C. One Primer was not Incorporated into Full Length  
15 Product

One of our most important findings is that the primer of the same polarity as the major RNA species was not detectably incorporated into full length DNA product. To demonstrate this, individual primers were terminally  
20 labeled with  $^{32}\text{P}$ , and the incorporation of each primer into DNA products was examined by polyacrylamide gel electrophoresis. We initially expected both primers to be incorporated into full length DNA products. However, the primer containing the promoter was not observed in full  
25 length DNA product. In fact, it remained quite small. These results were confirmed with a number of different targets and primer sets. Our method is explained below.

To identify the species of cDNA accumulated during autocatalysis, primers were  $^{32}\text{P}$ -labeled at the 5' end with  
30 T4 polynucleotide kinase and spiked into amplification reactions. The following examples show that cDNA of one polarity is preferentially accumulated during amplification, and that the accumulated strand is always complementary to the predominant RNA species synthesized  
35 during autocatalysis. Figure 5 shows the result of incorporation of  $^{32}\text{P}$ -labeled HIV region 2 primers during amplification. This primer set results in synthesis of a

214 base RNA of the (+) sense. The primer complementary to the RNA was incorporated into two major bands of 235 and 214 bases when target sequences were present (lane 3). No full length fragments were seen in the absence of target (lane 4). The 214 base fragment represents cDNA equal in length to the RNA while the 235 base fragment is equal in length to the RNA + 21 bases of the T7 promoter sequence. In contrast, the promoter primer was not observed in full length DNA product in the presence or absence of target (lanes 1 and 2 respectively).

Lanes 5-8 of Figure 5 show the result of incorporation of <sup>32</sup>P-labeled HBV region 1 primers during amplification. These are known as T7pro<sup>+</sup> and T7pro<sup>-</sup>. This primer set is capable of producing 132 base RNA of two polarities but the (+) strand RNA predominates. T7pro<sup>-</sup>, which is complementary to the predominant RNA, was incorporated into fragments of 132 bases and 153 bases consistent with our proposed mechanism (lane 7 (+) target, lane 8, (-) target). The 153 base fragment is equal in length to the RNA + 21 bases of the T7 promoter sequence of the T7 pro<sup>+</sup>. In contrast, <sup>32</sup>P-labeled T7pro<sup>+</sup> primer was not incorporated into fragments of either length (lane 5 (+) target, lane 6, (-) target).

The reactions analyzed by gel electrophoresis were also analyzed by HPA to determine if cDNA of the same polarity as the predominant RNA could be detected by another method. Plus and minus strand probes were used to determine the relative ratio of strands made, and a portion of each reaction was treated with RNase A to determine the ratio of RNA to DNA. The results are set forth below.

Table 23

	Probe Polarity	Prim r set	RLU	RLU
			No treatment	RNAs A
5	(-)	*HIV Region 2	679,182	1037
		HIV Region 2	453,675	1464
	(+)	*HIV Region 2	32,922	1,094,249
		HIV Region 2	39,494	655,595
	(-)	*HBV Region 1	567,110	4,854
10		HBV Region 1	671,656	4,210
	(+)	*HBV Region 1	56,550	303,160
		HBV Region 1	77,983	450,332

\* = (+) strand primer labeled with  $^{32}\text{P}$ , others, (-) strand primer labeled with  $^{32}\text{P}$ .

15

These results show that the amplifications worked well, even when full length product was not observed with the promoter primer. These results correlate with what was observed in the previous study, that is, most of the signal observed with one sense probe is from RNA, and the complementary strand signal is as expected, from DNA. This was true even for the HBV region 1 primer set which should have made RNA of both polarities.

#### 25 D. Confirmation of Mechanism Showing that the Enzyme Cuts RNA of the RNA/DNA Hybrid at Specific Loci

Based upon the experiments and observations reported hereinabove, the mechanism for the amplification systems that takes into account probable sequence specificities in the RNase H is depicted in Fig. 4. In support of this mechanism, since RNase H sequence specificity is a key element, it is necessary to show that indeed the enzymes cut the RNA of an RNA:DNA hybrid at specific locations. Furthermore, the locations of the cut sites needed to be in specific regions according to the model in order for good amplification to be obtained. To examine this question, an RNA:DNA hybrid was prepared that contained

the RNA that would be generated from a known target and primer set combination. The RNA was labeled with  $^{32}\text{P}$ , incubated with AMV reverse transcriptase (containing its associated RNase H activity), and the reaction products  
5 were analyzed by polycryamide gel electrophoresis. The results were entirely consistent with the new reaction mechanism, namely, the fragment size indicated that several small pieces were generated and that these were produced by cuts near one or both ends. The interior  
10 region of the molecule was not cut. This experiment confirmed that the enzyme has sequence or structural specificity under the reaction conditions used.

Further experiments were performed to determine where the cuts occurred since the proposed mechanism requires  
15 that multiple cuts occur in the region binding the promoter-containing primer. By labeling the termini of the RNA individually and analyzing the digestion products, it was demonstrated that the cuts were made only at the 5' end of the RNA. This also is consistent with the proposed  
20 mechanism.

#### Example 16

Figure 6 shows that the RNase H activities from AMV, MMLV and E. coli cut at specific RNase H cleavage sites.  
25 The arrows in the figure indicate the position of full-length RNA.

Figure 6 shows the result of an experiment in which HIV region 2 RNA was internally labelled with  $^{32}\text{P}$ , hybridized to a single-stranded target sequence and nicked  
30 with RNase H from AMV for 45 minutes (lane 2), MMLV for 5, 15, 45 or 120 minutes (lane 3-6) or E. coli RNase H for 5 minutes (lane 7). The sizes of fragments produced were discrete and different with each enzyme, indicating specificity of cleavage of the enzymes and varied  
35 specificity among enzymes with this template. The most rapid degradation was observed in the presence of E. coli

RNAse H, indicating less specificity or greater activity of cutting with this enzyme.

Figure 6b shows the results of hybridization of HBV region 1 RNA to a synthetic target sequence, followed by  
5 nicking with RNAse H from AMV reverse transcriptase for 5, 30, 60 or 120 minutes (lanes 2-5) or *E. coli* for 1, 3, 15 or 60 minutes (lanes 6-9). Different sized fragments were produced with the two enzymes, indicating specificity of cleavage. When the HBV RNA was labeled on the 3' terminus  
10 and nicked with AMV reverse transcriptase, the same sized fragments were observed, indicating that the cleavage sites were near the 5' end of the RNA.

These data indicate that specific sites are cleaved with RNAse H from AMV, MMLV and *E. coli*, and that at least  
15 some sites are different with the three enzymes. The presence of specific sites within the region to be amplified allows the RNA in an RNA:DNA hybrid to be cleaved, allowing autocatalysis to occur efficiently. Primers designed using cut site information show improved  
20 amplification efficiency. This is consistent with our observations that certain primer sets amplified to different extents depending on the source of RNAse H.

#### E. Identification of MMLV and AMV RNAse H cut sites.

##### 25 Example 17

To identify sites digested by AMV RNAse H, RNA was hybridized with a complementary sequence and nicked with AMV RNAse H. Following denaturation, a primer complementary to the 3' end of the region to be sequenced  
30 was hybridized and extended in the presence of dideoxynucleotides by the Sanger sequencing method. Termination of cDNA synthesis, indicating cleavage of RNA, was observed at the following sites for the HBV RNA:

5' GGGAGAGGUUAUCGC\*UGGA\*UGUGUCUGCGGCGUUUUAUCA\*UAU  
35 UCCUCUUCA\*UCCUG...3'

To identify sites digested by MMLV RNAse H, RNA was hybridized with a complementary sequence and nicked with

MMLV RNase H. Following denaturation, a primer complementary to the 3' end of the region to be sequenced was hybridized and extended in the presence of dideoxynucleotides by the Sanger sequencing method.

- 5 Termination of cDNA synthesis was observed at the following sites for the HBV RNA:

5' GGGAGAGGUUAUCGC\*UGGA\*UGUGUCUGCGGC\*GUUUUAUCA\*  
UAUCCUCUUAUCCUGC\*UGCUAUGCCUCA\*UCUUC...-3'

- 10 The following sites were identified for a second HBV RNA sequence:

5' GGGAGACCCGAGAU\*UGA\*GAUCUUCUGCGAC  
GCGGCGAU\*UGA\*GAUCUGCGUCU\*GCGAGGCGAGGGAGU\*UCU\*UCUU\*CUA  
GGGGACCUGCCUCGGUCCCGUC\*GUCUA...3'

- 15 The following sites were identified for an HIV RNA sequence:

5' GGGAGACAAA\*UGGCAGUA\*UUCAUCCAC  
AAUUUAAAAGAAAAGGGGGGAUUGGGGGGUA  
CAGUGCAGGGGAAAGAAUAGUAGACAUAAUAGC\*AAACAGACA  
UAC\*AAACUAAAGAAUACAAAAACAAUAC\*AAAAAUCAA  
20 AAUUUUCGGGUUUAUUACAGGGAC\*AGC\*AGAAA...3'

- Most of the cleavage sites occurred near the dinucleotides CA or UG. The method used for detecting cleavage sites only identified sites which accumulated during the cleavage reaction. It is expected that additional sites  
25 could be cleaved which were not recognized by the method used.

#### F. Primers for Amplification Systems

- Based on findings that the various RNase H enzymes  
30 have sequence specificity, we have tested various primer/target combinations and attempted to optimize their performance in amplification systems. Data obtained to date indicates that the piece size of the RNA fragments produced is relatively large and that the fragments  
35 probably do not spontaneously dissociate from the duplex. This is not unexpected since work with AMV reverse transcriptase copying AMV RNA or poliovirus RNA showed that

the RNA fragments that were produced by the RNase H were used by the enzyme to prime the synthesis of cDNA from the initially synthesized cDNA strand.

If the RNase H enzymes have sequence specificity, the  
5 amplification reaction proceeds as follows (beginning with the RNA intermediate in the reaction):

The primer complementary to the major RNA species produced during amplification binds at the 3' terminus of the RNA. Since the concentration of primer is high in the  
10 reaction, excess primer produces RNA:DNA duplexes which may be cut by the RNase H activity before being able to initiate synthesis. Therefore, it is preferable that the primer binding region does not contain a large number of sequences recognized by the RNase H enzyme used in the  
15 reaction.

As cut sites occur frequently, it may not be practical in some cases to design an RNA complementary primer without recognized cut sites; in such cases, the cut sites should be as near the 5' terminus as possible to  
20 allow the 3' terminal portion of the primer to remain annealed to the RNA.

Upon extension of the primer by a suitable DNA polymerase, the binding site for the second primer, which contains the RNA polymerase promoter, must now be exposed.  
25 It is sufficient to remove only a portion of the RNA to allow nucleation and zippering of the primer as it hybridizes to the cDNA, to allow reverse transcriptase mediated binding of the primer and initiation of synthesis, or merely to nick the RNA so that the RNA  
30 fragment that results may be displaced. Since our data show relatively large pieces of RNA are made and that the promoter containing primer is not incorporated into full-length DNA, the following events can occur:

1. There is sufficient nicking of the RNA to permit  
35 binding of the promoter-primer. Whether a nick in the appropriate place simply produces an RNA fragment sufficiently small to melt off and expose the primer

binding site or a portion thereof or whether a nick allows an unwinding activity associated with one or more of the enzymes to displace the RNA fragment is not known at this time.

5        2.    The cDNA 3' terminus is extended by the reverse transcriptase to make the promoter region double-stranded DNA.

         3.    RNA is synthesized from the complex thus made. This complex would consist of a cDNA bound to RNA and  
10 containing a double-stranded DNA promoter.

         Thus, there must be a sequence recognized by the RNase H activity present in the reaction somewhere in or near the binding site for the primer containing the RNA polymerase promoter.

15        In some applications, it may also be desirable to not have RNase H recognition sites within the target sequence itself. Sites within the target may be cleaved and used to produce RNA primers for synthesis of double-stranded cDNA regions. It may be preferable to eliminate the  
20 possibility of this enzymatic activity.

         New primer sets were designed based upon the model and the RNase H sequence specificity information that we have obtained. Our design criteria are as follows:

For the T7 promoter-primer:

- 25    1)    The primer region should have one or more cut sites per 10 bases.
- 2)    The primer region should have a cut site near the 5' end.
- 3)    The primer region should have a cut site near the 3'  
30 end and possibly a partial site at the 3' end.
- 4)    The primer length should be >18 bases.
- 5)    The  $T_m$  estimated should be about 55-65°C.

For the other primer:

- 1)    The primer should have few or no RNase H cut sites.
- 35    2)    Any cut sites in the primer should be near the 5' end.

3) The primer length should be about 18-28 bases in length.

4) The  $T_m$  estimated should be about 55-65°C.

Significantly better synthesis was obtained from primer sets designed using these criteria and knowledge of the mechanism and sequence specificities. This shows the utility of the invention in making possible the design of functional primer sets for specific target regions. These are explained more fully below.

10

#### Example 18

Our findings regarding RNase H specificity have been used to design efficient promoter-primer combinations. Prior art methods simply nonselectively attached promoters to primer sequences. We have been able to design and optimize promoter-primer combinations to increase the yield of amplified product. The following experiment shows that small changes in promoter-primer sequence result in large changes in amplification efficiency.

20 The following examples show primers from similar regions which were compared for RNase H cleavage sites and GP-III amplification efficiency. In each example, duplicate amplifications were performed using common reagents except for the primers being tested.

#### 25 1. non-promoter primers.

In the first example, the non-promoter primer site for the CML major t(14; 18) breakpoint amplification region was moved 15 bases, resulting in a reduction in the number of putative RNase H cut sites from 4 to 1, assuming a 4 base recognition sequence or from 5 to 2 assuming a 3 base recognition sequence. The reaction was performed as described for Example 15 except that 2.5 mM ATP, 16.5 mM MgCl<sub>2</sub> and 50 mM KCl were included. This change in primer sequence had a dramatic positive effect on amplification efficiency. In the second case, an intentional mismatch was placed internally in the non promoter primer of HBV region 1 to remove the only putative RNase H cut site,

assuming a 4 base recognition site. In the case of a 3 base cut site, one skilled in the art would recognize that the mismatch removed the cut site nearest the 3' end. This change also had a definitive positive effect on amplification efficiency. The data demonstrate that two methods, changing the position of the primer, or inclusion of mismatches, can be used to enhance amplification. Presumably, removal of RNase H cut sites from the non-promoter primer results in more efficient priming of cDNA synthesis during autocatalysis.

	Sequence	RLU
Example 1	GGAGCTGCAGATGCTGACCAAC	78,880
	GACCAACTCGTGTGTGAAACTCCA	2,552,333
Example 2	TCCTGGAATTAGAGGACAAACGGGC	57,710
	TCCTGGAATTAGAGGATAAACGGGC	518,695

2. promoter-primers. The following examples show promoter primers which come from similar regions but which differ in the number of putative RNase H cut sites. In the first case, the two promoter primer sites for the HIV region 5 are displaced by 10 bases, changing the number of putative RNase H cut sites from two to three, assuming a four base recognition site, or from 3 to 5 assuming a 3 base recognition site. This has a positive effect on amplification efficiency. In the second case, a sequence containing putative RNase H cut sites was inserted upstream of the promoter primer for the major breakpoint t(14); 18) translocation, and one mismatch to the target was included to generate a cut site within the primer region. This also had a positive effect on amplification efficiency. This demonstrates that insertion of RNase H cut sites in the promoter primer can be used to enhance amplification efficiency. Presumably, inclusion of RNase H cut sites assists in RNA strand displacement, increasing the efficiency of copying of the promoter region, thereby resulting in more efficient autocatalysis.

	Primer name	RLU
Exempl 3	A	45,466
	B	908,147
Example 4	C	64,601
5	D	2,946,706

Sequences of the primers above are:

Primer A:

AATTTTAATACGACTCACTATAGGGAGAAATCTTGTGGGGTGGCTCCTTCT-3'

Primer B:

10 AATTTAATACGACTCACTATAGGGAGAGGGGTGGCTCCTTCTGATAA TGCTG-3'

Primer C:

ATTTAATACGACTCACTATAGGGAGACGGTGACCGTGGTCCCTTG-3'

Primer D:

TAAATTAATACGACTCACTATAGGGAGATCAGTTACAATCGC

15 TGGTATCAACGCTGAGCAGACGCTGACCGTGGTCCCTTG-3'

In the above examples, removal of RNase H cut sites from the non-promoter primer resulted in enhanced amplification, even if the removal of the cut site involved the incorporation of a mismatch to the original target. Design of the promoter-containing primer to include additional RNase H cut sites also enhanced amplification, again, even if the incorporation of cut sites involved inclusion of mismatches to the original target. The number, distribution, and position of putative RNase H cut sites determine, in part, the usefulness of a given primer.

Improvement of amplification by inclusion of intentional mismatches or insertion of sequences between the promoter and primer are nonobvious improvements to the amplification method.

In a preferred embodiment of the present invention, the RNA target sequence is determined and then analyzed to determine where RNase H degradation will cause cuts or removal of sections of RNA from the duplex. Experiments can be conducted to determine the effect of the RNase degradation of the target sequence by RNase H present in AMV reverse transcriptase and MMLV reverse

transcriptase, by E. coli RNase H or by combinations thereof.

In selecting a primer, it is preferable that the primer be selected so that it will hybridize to a section  
5 of RNA which is substantially nondegraded by the RNase H present in the reaction mixture. If there is substantial degradation, the cuts in the RNA strand in the region of the primer may stop or inhibit DNA synthesis and prevent extension of the primer. Thus, it is desirable to select  
10 a primer which will hybridize with a sequence of the RNA target, located so that when the RNA is subjected to RNase H, there is no substantial degradation which would prevent formation of the primer extension product.

The site for hybridization of the promoter-primer is  
15 chosen so that sufficient degradation of the RNA strand occurs to permit removal of the portion of the RNA strand hybridized to the portion of the DNA strand to which the promoter-primer will hybridize. Typically, only portions of RNA are removed from the RNA:DNA duplex by RNase H  
20 degradation and a substantial part of the RNA strand remains in the duplex. An RNA:DNA duplex containing a double-stranded DNA promoter results.

Claims:

1. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:
  - (a) treating a nucleic acid which comprises an  
5 RNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith and which optionally has a sequence 5' to the complexing sequence  
10 which includes a promoter for an RNA polymerase, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;
  - (b) extending the primer in an extension reaction using the target as a template to give a DNA  
15 primer extension product complementary to the RNA target;
  - (c) separating the DNA primer extension product from the RNA target using an enzyme which selectively degrades the RNA target;
  - (d) treating the DNA primer extension product  
20 with a second oligonucleotide which comprises a primer or a splice template and which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith, under conditions whereby an oligonucleotide/target sequence complex is  
25 formed and DNA synthesis may be initiated, provided that if the first oligonucleotide does not have a promoter, then the second oligonucleotide is a splice template which has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase;
  - 30 (e) extending the 3'-terminus of either the second oligonucleotide or the first primer extension product, or both, in a DNA extension reaction to produce a template for an RNA polymerase; and
  - (f) using the template of step (e) to produce  
35 multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter sequence.

2. A method according to claim 1 wherein the enzyme of step (c) comprises an RNase H.

3. A method according to claim 1 further comprising using the oligonucleotides and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

4. A method according to claim 3 wherein the enzyme of step (c) comprises an RNase H.

5. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) treating a target nucleic acid which comprises an RNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;

(b) extending the first primer in an extension reaction using the target as a template to give a first DNA primer extension product complementary to the RNA target;

(c) separating the first DNA primer extension product from the RNA target using an enzyme which selectively degrades the RNA target;

(d) treating the first DNA primer extension product with a second oligonucleotide which comprises a second primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith, under conditions whereby an oligonucleotide/target sequence is formed and DNA synthesis may be initiated;

( ) extending th 3'-terminus of the second primer in a DNA extension reacti n to give a second DNA primer extension product, thereby producing a template for an RNA polymerase; and

- 5 (f) using the template of step (e) to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter sequence.

6. A method according to claim 5 wherein said  
10 enzyme of step (c) comprises an RNase H.

7. A method according to claim 5 wherein the second primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

15

8. A method according to claim 7 wherein the enzyme of step (c) comprises an RNase H.

9. A method according to claim 5 further compris-  
20 ing using the primers and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

10. A method according to claim 9 wherein the second primer has a sequence 5' to the complexing sequence  
25 which includes a promoter for an RNA polymerase.

11. A method according to claim 5 further comprising:

(g) treating an RNA copy from step (f) with  
30 the second primer under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated.

(h) extending the 3'-terminus of the second primer in a DNA extension reaction to give a second DNA  
35 primer extension product which is complementary to the RNA copy;

86

(i) separating the second DNA primer extension product from the RNA copy using an enzyme which selectively degrades the RNA copy;

(j) treating the second DNA primer extension product with the first primer under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated.

(k) extending the 3'-terminus of the first primer in a DNA extension reaction to give a first DNA primer extension product and the 3'-terminus of the second DNA primer extension product, thereby producing a template for a RNA polymerase; and

(l) using the template of step (k) to produce multiple copies of the target sequence using an RNA polymerase which recognizes the promoter.

12. A method according to claim 11 wherein the second primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

13. A method according to claim 11 further comprising using the primers and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

14. A method according to claim 13 wherein the second primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

15. A method according to claim 11 further comprising:

(m) using the RNA copies of step (l), repeating steps (g) to (l) to autocatalytically synthesize multiple copies of the target sequence.

16. A method according to claim 15 wherein the enzyme of steps (c) and (i) comprises an RNase H.

17. A method according to claim 15 wherein the second primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

5 18. A method according to claim 17 wherein the enzyme of steps (c) and (i) comprises an RNase H.

19. A method according to claim 7 further comprising:

10 (g) treating the RNA copies of step (f) with the first and second primers under complexing conditions;

(h) extending the primers in a DNA extension reaction using the RNA copies as templates to give DNA primer extension products;

15 (i) separating the DNA primer extension products from the RNA copies using an enzyme which selectively degrades the RNA copies;

(j) treating the DNA primer extension products with the primers under complexing conditions;

20 (k) extending the primers in a DNA extension reaction to give a complementary primer extension product, thereby producing templates for a RNA polymerase; and

(l) using the templates of step (k) to produce multiple copies of the target sequence using an RNA  
25 polymerase which recognizes the promoter.

20. A method according to claim 19 further comprising using the primers and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

30

21. A method according to claim 19 further comprising:

(m) using the RNA copies of step (l), repeating steps (g) to (l) to autocatalytically synthesize  
35 multiple copies of the target sequence.

22. A method according to claim 21 wherein the enzyme of steps (c) and (i) comprises an RNase H.

23. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) treating a nucleic acid which comprises an RNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;

(b) extending the 3' terminus of the first primer in an extension reaction using the target as a template to give a DNA primer extension product complementary to the RNA target;

(c) separating the DNA primer extension product from the RNA target using an enzyme which selectively degrades the RNA target;

(d) treating the DNA primer extension product with a second oligonucleotide which comprises a splice template which has a complexing sequence sufficiently complementary to the 3' terminus of the primer extension product to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase, under conditions whereby an oligonucleotide/-target sequence complex is formed and DNA synthesis may be initiated;

(e) extending the 3'-terminus of the primer extension product to add thereto a sequence complementary to the promoter, thereby producing a template for an RNA polymerase;

(f) using the template of step (e) to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter.

24. A method according to claim 23 further comprising using the oligonucleotides and the RNA copies to autocatalytically synthesize multiple copies of the target sequence.

5

25. A method according to claim 24 wherein the 3' terminus of the splice template is blocked.

26. A method according to claim 23 further comprising:

(g) using the RNA copies of step (f), repeating steps (a) to (f) to autocatalytically synthesize multiple copies of the target sequence.

27. A method according to claim 26 wherein said enzyme of step (c) comprises an RNase H.

28. A method according to claim 26 wherein the 3' terminus of the splice template is blocked.

20

29. A method according to claim 26 wherein in step (e), the splice template acts as a second primer and the 3' terminus of the splice template is extended in an DNA extension reaction to give a second primer extension product which is complementary to the first primer extension product.

30. A method according to claim 29 wherein the first primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

31. A method according to claim 30 wherein said enzyme of step (c) comprises an RNase H.

32. A method according to claim 28 wherein said primer and said splice template comprise a single olig nucleotide having a sequence which comprises the

splice template 5' to a sequence which comprises the primer.

33. A method of synthesizing multiple copies of a  
5 target sequence which comprises:

(a) treating a single stranded target nucleic acid which comprises a DNA target sequence having a defined 3' terminus with a first oligonucleotide which comprises a splice template which has a complexing  
10 sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase, under conditions whereby an oligonucleotide/target sequence complex is formed and  
15 DNA synthesis may be initiated;

(b) extending the 3' terminus of the target to add a sequence complementary to the promoter, thereby producing a template for an RNA polymerase; and

(c) using the template of step (b) to produce  
20 multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter.

34. A method according to claim 33 wherein the 3' terminus of the splice template is blocked.  
25

35. A method according to claim 33 further comprising:

(d) treating an RNA copy of step (c) with a second oligonucleotide which comprises a primer which has  
30 a complexing sequence sufficiently complementary to the 3' terminal portion of the RNA copy to complex therewith, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;

35 (e) extending the 3' terminus of the primer in a DNA extension reaction to give a DNA primer extension product which is complementary to the RNA copy;

(f) separating the DNA primer extension product from the RNA copy;

(g) treating the DNA primer extension product with the splice template under conditions whereby an  
5 oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;

(h) extending the 3' terminus of the primer extension product in a DNA extension reaction to add a sequence complementary to the promoter, thereby producing  
10 a template for an RNA polymerase; and

(i) using the template of step (h) to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter.

15 36. A method according to claim 35 wherein step (f) uses an enzyme which selectively degrades the RNA copy.

37. A method according to claim 36 further comprising using the oligonucleotides and the RNA copies  
20 of step (i), to autocatalytically synthesize multiple copies of the target sequence.

38. A method according to claim 37 wherein the 3' terminus of the splice template is blocked.

25

39. A method according to claim 36 further comprising:

(j) using the RNA copies of step (i), repeating steps (d) to (i) to autocatalytically synthesize  
30 multiple copies of the target sequence.

40. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) treating a single stranded target nucleic  
35 acid which comprises a DNA target sequence with a first oligonucleotide which comprises a first primer which has a complexing sequence sufficiently complementary to the

3'-terminal portion of the target sequence to complex  
th rewith and a sequence which includes a promoter for an  
RNA polymerase, under conditions whereby an  
oligonucleotide/target sequence complex is formed and DNA  
5 synthesis may be initiated;

(b) extending the primer in a DNA extension  
reaction using the target as a template to give a first  
DNA primer extension product complementary to the DNA  
target sequence;

10 (c) separating the first primer extension  
product from the target;

(d) treating the first primer extension  
product with a second oligonucleotide which comprises a  
second primer which has a complexing sequence sufficiently  
15 complementary to the 3'-terminal portion of the target  
sequence to complex therewith, under conditions whereby an  
oligonucleotide/target sequence complex is formed and DNA  
synthesis may be initiated;

(e) extending the 3' terminus of the second  
20 primer in a DNA extension reaction to give a second DNA  
primer extension product, thereby producing a template for  
a RNA polymerase;

(f) using the template of step (e) to produce  
multiple RNA copies of the target sequence using an RNA  
25 polymerase which recognizes the promoter;

(g) treating an RNA copy from step (f) with  
the second primer under conditions whereby an  
oligonucleotide/target sequence complex is formed and DNA  
synthesis may be initiated;

30 (h) extending the 3'-terminus of the second  
primer in a DNA extension reaction to give a second DNA  
primer extension product complementary to the RNA copy;

(i) separating the DNA primer extension  
product from the RNA copy using an enzyme which  
35 selectively degrades the RNA copy;

(j) treating the second primer extension  
product with the first primer, under conditi ns wher by an

oligonucleotide/target complex is formed and DNA synthesis may be initiated;

(k) extending the 3'-terminus of the first primer in a DNA extension reaction to give a first DNA primer extension product and the 3'-terminus of the second primer extension product, thereby producing a template for an RNA polymerase; and

(l) using the template of step (k) to produce multiple copies of the target sequence using an RNA polymerase which recognizes the promoter.

41. A method according to claim 40 wherein the enzyme of step (i) comprises an RNase H.

42. A method according to claim 40 wherein the second primer has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

43. A method according to claim 42 wherein the enzyme of step (i) comprises an RNase H.

44. A method according to claim 42 further comprising using the oligonucleotides and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

45. A method according to claim 44 wherein the enzyme of step (i) comprises an RNase H.

46. A method according to claim 42 further comprising:

(m) using the RNA copies of step (l) repeating steps (g) to (l) to autocatalytically synthesize multiple copies of the target sequence.

47. A method according to claim 46 wherein the enzyme of step (i) comprises an RNase H.

48. A method of synth sizing multiple copies of a target nucleic acid sequence which comprises:

(a) treating a single stranded target nucleic acid which comprises a DNA target sequence with a first  
5 oligonucleotide which comprises a primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence to complex therewith under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be  
10 initiated;

(b) extending the 3'-terminus of the primer in an extension reaction using the target as a template to give a DNA primer extension product complementary to the target;

15 (c) separating the primer extension product from the target;

(d) treating the primer extension product with a second oligonucleotide which comprises a splice template which has a complexing sequence sufficiently complementary  
20 to the 3' terminal portion of the target sequence to complex therewith and a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase, under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be  
25 initiated;

(e) extending the 3'-terminus of the primer extension product to add thereto a sequence complementary to the promoter, thereby producing a template for the RNA polymerase; and

30 (f) using the template of step (e) to produce multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter;

49. A method according to claim 48 further  
35 comprising:

(g) treating an RNA copy of step (f) with the primer under conditions whereby an oligonucleotide/target

95

sequence complex is formed and DNA synthesis may be initiated;

(h) extending the 3'-terminus of the primer in a DNA extension reaction to give a primer extension  
5 product;

(i) separating the primer extension from the RNA copy using an enzyme which selectively degrades the RNA copy;

(j) treating the primer extension product with  
10 the splice template under conditions whereby an oligonucleotide/target sequence complex is formed and DNA synthesis may be initiated;

(k) extending the 3' terminus of the primer extension product to add thereto a sequence complementary  
15 to the promoter, thereby producing a template for an RNA polymerase;

(l) using the template of step (k) to produce RNA copies of the target sequence using an RNA polymerase which recognizes the promoter.  
20

50. A method according to claim 49 wherein the 3' terminus of the splice template is blocked.

51. A method according to claim 49 further  
25 comprising using the oligonucleotides and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

52. A method according to claim 51 wherein the  
30 enzyme of step (i) is an RNase H.

53. A method according to claim 49 further comprising:

(m) using the RNA copies of step (l),  
35 repeating steps (g) to (l) to autocatalytically synthesize multiple copies of the target sequence.

54. A method according to claim 53 wherein the enzym of step (i) is an RNase H.

55. A method for autocatalytically synthesizing  
5 multiple copies of a target nucleic acid sequence under  
conditions of substantially constant temperature, ionic  
strength and pH which comprises:
- (a) combining
    - (1) a target nucleic acid which comprises  
10 a single-stranded RNA target sequence;
    - (2) a first oligonucleotide which  
comprises a primer which has a complexing sequence  
sufficiently complementary to the 3' terminal portion of  
the RNA target sequence to complex therewith and which  
15 optionally has a sequence 5' to the complexing sequence  
which includes a promoter for an RNA polymerase;
    - (3) a second oligonucleotide which  
comprises a primer or a splice template and which has a  
complexing sequence sufficiently complementary to the 3'-  
20 terminal portion of a complement to the RNA target  
sequence to complex therewith, provided that if the first  
oligonucleotide does not have a promoter, then the second  
oligonucleotide is a splice template which has a sequence  
5' to the complexing sequence which includes a promoter  
25 for an RNA polymerase;
    - (4) a DNA dependent DNA polymerase;
    - (5) an RNA dependent DNA polymerase;
    - (6) an enzyme which selectively degrades  
the RNA strand of an RNA:DNA duplex; and
    - 30 (7) an RNA polymerase which recognizes  
the promoter; and
  - (b) incubating the mixture of step (a) under  
DNA priming and nucleic acid synthesizing conditions,  
which include substantially constant temp rature, ionic  
35 strength and pH.

56. A method according to claim 55 wherein the enzyme of step (a)(6) comprises an RNase H.

57. A method according to claim 55 wherein the first primer has a promoter.

58. A method according to claim 57 wherein the second oligonucleotide comprises a second primer which has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase.

59. A method according to claim 58 wherein the DNA dependent and RNA dependent DNA polymerase activities and RNase H activities comprise a reverse transcriptase.

15

60. A method for autocatalytically synthesizing multiple copies of a target nucleic acid sequence under conditions of substantially constant temperature, ionic strength and pH which comprises:

20

(a) combining:

(1) a target nucleic acid which comprises an RNA target sequence;

(2) a primer and a splice template of opposite sense wherein one has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence of the RNA target to complex therewith and the other has a complexing sequence sufficiently complementary to the 3' terminal portion of the complement of the target sequence to complex therewith and wherein the splice template has a sequence 5' to the complexing sequence which includes a promoter;

(3) a DNA dependent DNA polymerase;

(4) an RNA dependent DNA polymerase;

(5) an enzyme which selectively degrades the RNA strand of an RNA:DNA duplex; and

35

(6) an RNA polymerase which recognizes the promoter of the splice template; and

(b) incubating the mixture of step (a) under DNA priming and nucleic acid synthesizing conditions which include substantially constant temperature, ionic strength, and pH.

5

61. A method for autocatalytically synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) combining:

10 (1) a nucleic acid which comprises a single stranded DNA target sequence.

(2) a first primer which has a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex therewith and a  
15 sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase;

(3) a DNA polymerase;

(b) incubating the mixture of step (a) under DNA priming and synthesizing conditions whereby a primer  
20 extension product complementary to the target sequence is synthesized using the target sequence as a template;

(c) treating the reaction mixture of step (b) to cause separation of DNA duplexes

(d) adding to the reaction mixture of step  
25 (c):

(1) a second primer which has a complexing sequence sufficiently complementary to the 3'-terminal portion of the target sequence of the primer extension product to complex therewith.

30 (2) a DNA dependent DNA polymerase;

(3) an RNA dependent DNA polymerase;

(4) an enzyme which selectively degrades the RNA strand of an RNA:DNA complex; and

(5) an RNA polymerase which recognizes  
35 the promoter; and

(e) incubating the mixture of step (d) under DNA priming and nucleic acid synthesizing conditions which

include substantially constant temperature, ionic strength and pH.

62. A method for autocatalytically synthesizing multiple copies of a target nucleic sequence under conditions of substantially constant temperature, ionic strength and pH which comprises:

(a) combining

(1) a target nucleic acid which comprises a target sequence;

(2) a first oligonucleotide which comprises first primer which has a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex thereto and which optionally has a sequence 5' to the complexing sequence which includes a promoter for an RNA polymerase;

(3) a second oligonucleotide which comprises a second primer or a splice template and which has a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence to complex thereto, provided that if the first oligonucleotide does not have a promoter, then the second oligonucleotide is a splice template, which has a sequence 5' to the complexing sequence which includes a promoter;

(4) a DNA dependent DNA polymerase;

(5) an RNA dependent DNA polymerase;

(6) an enzyme which selectively degrades the RNA strand of an RNA: DNA duplex; and

(7) an RNA polymerase; and

(b) incubating the mixture of step (a) under DNA priming and DNA synthesizing conditions, which include substantially constant temperature, ionic strength and pH.

63. A method according to claim 61 wherein the first olig nucleotide has a promoter sequence.

64. A method according to claim 62 wherein the enzyme of step (a)(6) comprises an RNase H.

5 65. A method according to claim 63 wherein the enzyme of step (a)(6) comprises an RNase H activity present in the reverse transcriptase.

66. A method according to claim 64 wherein the enzyme of step (a)(5) comprises added exogenous RNase H.  
10

67. A method according to claim 65 wherein the second oligonucleotide has a promoter.

68. A method according to claim 61 wherein the enzyme of step (a)(6) comprises an RNase H.  
15

69. A method according to claim 67 wherein the enzyme of step (a)(6) comprises an RNase H activity in the reverse transcriptase.  
20

70. A method according to claim 67 wherein the enzyme of step (a)(6) comprises added exogenous RNase H.

71. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:  
25

(a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with the said portion of the RNA target, said portion of the RNA target located such that it remains substantially undegraded;  
30

(b) selecting a promoter-primer, complementary a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where substantially all of the complementary RNA is removed from the duplex by degradation of said RNA; and,  
35

(c) combining said RNA target with said promoter-primer, reverse transcriptase having associated

RNAse H activity, and transcriptas and forming multiple copies of said target nucleic acid s quenc .

72. A method according to claim 71 wherein  
5 exogenous RNAse H is added at step (c).

73. A method according to claim 72 wherein the reverse transcriptase of step (c) lacks associated RNAse H activity.

10

74. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with  
15 the said portion of the RNA target, said portion of the RNA target located such that it remains capable of forming a primer extension product after being exposed to degradation by a selected RNAse H;

(b) selecting a promoter-primer, complementary  
20 a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where substantially all of the complementary RNA is removed from the duplex by degradation of said RNA; and

(c) combining said RNA target with said  
25 promoter-primer, reverse transcriptase having associated RNAse H activity and transcriptase and forming multiple copies of said target nucleic acid sequence.

75. A method according to claim 74 wherein  
30 exogenous RNAse H is added at step (c).

76. A method according to claim 75 wherein the reverse transcriptase of step (c) lacks associated RNAse H activity.

35

77. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with the said portion of the RNA target, said portion of the RNA target located such that it remains substantially  
5 undegraded;

(b) selecting a promoter-primer, complementary to a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where substantially all of the complementary RNA is removed from  
10 the duplex by degradation of said RNA; and,

(c) combining said RNA target with said promoter-primer, reverse transcriptase having associated RNase H activity and transcriptase and forming multiple copies of said RNA and multiple copies of DNA complementary to said RNA without making a substantially equivalent  
15 number of copies of DNA of the same polarity as said RNA target sequence.

78. A method according to claim 77 wherein  
20 exogenous RNaseH is added at step (c).

79. A method according to claim 78 wherein the reverse transcriptase of step (c) lacks associated RNase H activity.  
25

80. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

(a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with  
30 the said portion of the RNA target, said portion of the RNA target located such that it remains capable of forming a primer extension product after being exposed to degradation by a selected RNase H;

(b) selecting a promoter-primer, complementary  
35 a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where

substantially all of the complementary RNA is removed from the duplex by degradation of said RNA; and

- (c) combining said RNA target with said promoter-primer, reverse transcriptase having associated  
5 RNase H activity and transcriptase and forming multiple copies of said RNA and multiple copies of DNA complementary to said RNA without making a substantially equivalent number of copies of DNA of the same polarity as said RNA target sequence.

10

81. A method according to claim 80 wherein exogenous RNase H is added at step (c).

82. A method according to claim 81 wherein the  
15 reverse transcriptase of step (c) lacks associated RNase H activity.

83. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

- 20 (a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with the said portion of the RNA target, said portion of the RNA target located such that it remains capable of forming a primer extension product after being exposed to  
25 degradation by a selected RNase H;

- (b) selecting a promoter-primer, complementary a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where a sufficient amount of the 5' end of said RNA is removed from the  
30 duplex so as to permit the further extension of the primer extension product DNA to produce a complement to said promoter-primer; and

- (c) combining said RNA target with said promoter-primer, reverse transcriptase having associated  
35 RNase H activity and transcriptase and forming multiple copies of said target nucleic acid sequence.

104

84. A method according to claim 83 wherein exogenous RNase H is added at step (c).

85. A method according to claim 84 wherein the reverse transcriptase of step (c) lacks associated RNase H activity.

86. A method of synthesizing multiple copies of a target nucleic acid sequence which comprises:

10 (a) selecting a primer, complementary to a portion of an RNA target sequence, which complexes with the said portion of the RNA target, said portion of the RNA target located such that it remains capable of forming a primer extension product after being exposed to  
15 degradation by a selected RNase H;

(b) selecting a promoter-primer, complementary to a portion of the DNA obtained by extension of said primer, which complexes with the DNA in an area where a sufficient amount of the 5' end of said RNA is removed so  
20 as to permit said promoter-primer to complex to the 3' end of said DNA obtained by extension of said primer,

(c) combining said RNA target with said promoter-primer, reverse transcriptase having associated RNase H activity and transcriptase and forming multiple  
25 copies of said target nucleic acid sequence.

87. A method according to claim 86 wherein exogenous RNase H is added at step (c).

30 88. A method according to claim 87 wherein the reverse transcriptase of step (c) lacks associated RNase H activity.

89. A method of synthesizing multiple copies of a  
35 target nucleic acid sequence which comprises:

(a) treating a nucleic acid which comprises an RNA target sequence with a first oligonucleotide which

c mprises a first primer which has a complexing sequ nce sufficiently complementary to th 3'-terminal portion of the target sequence to complex therewith under conditions whereby DNA synthesis may be initiated;

5 (b) extending the primer in an extension reaction using the target as a template to give a DNA primer extension product complementary to the RNA target;

(c) selectively digesting the 5' end of said RNA target;

10 (d) treating the DNA primer extension product with a second oligonucleotide which comprises a primer or a splice template and which has a complexing sequence sufficiently complementary to the 3'-terminal portion of said DNA primer extension product under conditions whereby  
15 DNA synthesis may be initiated,

(e) extending the 3'-terminus of the first primer extension product in a DNA extension reaction to produce a template for an RNA polymerase; and

(f) using the template of step (e) to produce  
20 multiple RNA copies of the target sequence using an RNA polymerase which recognizes the promoter sequence.


90. A method according to claim 89 wherein the degradation of step (c) is enzymatic.

25

91. A method according to claim 90 using RNase H for enzymatic degradation.


92. A method according to claim 80 further  
30 comprising using the oligonucleotides and RNA copies to autocatalytically synthesize multiple copies of the target sequence.

- 1 / 27



SUBSTITUTE SHEET

FIG. 1b.



- 3 / 27




FIG. 1c.

- 4 / 27





FIG. 1d.

SUBSTITUTE SHEET


- 5 / 27

FIG. 1e.



- 6 / 27


FIG. 1f.



SUBSTITUTE SHEET


- 7 / 27

FIG. 1g.



- 8 / 27


FIG. 1h.



GO TO FIGURE 1e, STEP 7.

SUBSTITUTE SHEET,

- 9 / 27



*FIG. 1i.*  
**SUBSTITUTE SHEET,**

10/27





FIG. 1j.  
SUBSTITUTE SHEET

11/27

FIG. 1k.



SUBSTITUTE SHEET

12 / 27




FIG. 11.

SUBSTITUTE SHEET

13 / 27





FIG. 1m.


14 / 27



SUBSTITUTE SHEET

15/27

FIG. 10.



SUBSTITUTE SHEET

16/27





FIG. 2a.

SUBSTITUTE SHEET,


17/27

FIG. 2b.



18 / 27


FIG. 2c.



SUBSTITUTE SHEET

19/27


FIG. 2d.



SUBSTITUTE SHEET

20/27


FIG. 2e.



SUBSTITUTE SHEET

21 / 27

FIG. 3.



SUBSTITUTE SHEET,

22/27

## AMPLIFICATION SYSTEM—AUTOCATALYTIC CYCLE




FIG. 4a.




FIG. 4b.

24/27

AMPLIFICATION SYSTEM-GENERAL METHOD  
RNA TARGET WITH NON-TARGET SEQUENCES 5' TO TARGET REGION  
INITIATION MECHANISM I




FIG. 4c.

## RNA TARGET WITH NON-TARGET SEQUENCES 5' TO TARGET REGION INITIATION MECHANISM II




FIG. 4d.



FIG. 5

27/27





FIG. 6

# INTERNATIONAL SEARCH REPORT

International Application No PCT/US90/03907

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>3</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5):C12Q 1/68; C12P 19/34; C07H 15/12; G01N 33/53 U.S. CL.: 435/6, 91; 536/27; 28		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>4</sup>		
Classification System	Classification Symbols	
U.S	435/6; 435/91; 536/27; 536/28	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>5</sup>		
APS, CAS, BIOSYS		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT</b> <sup>14</sup>		
Category <sup>6</sup>	Citation of Document, <sup>10</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No. <sup>18</sup>
Y	US, A, 4,683,195 (Mullis et al.) 28 July 1987. See example 9, column 29.	1,5,7,23,33,35 40,42,48,71,83, 86
Y	Science, Volume 239, issued 29 January 1988, Stoflet et al. "Genomic Amplification with Transcript Sequencing " pages 491-494, see abstract.	1,5,7,23,33,35, 40,42,48,71,83, 86
<u>X</u> Y	Proceedings National Academy Sciences USA, Volume 86, issued February 1989, Kwoh et al. "Transcription-based amplification system and detection of amplified human immunodeficiency virus type 1 with a bread-based sandwich hybridization format" pages 1173-1177, see abstract.	1,5,23,33,35,40 42,48,71,83,86,7 2-4,6,8-22,24- 32,34,36-39,41, 43-47,49-70,72- 82,84,85,87- 92
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>15</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search <sup>2</sup>		Date of Mailing of this International Search Report <sup>2</sup>
10 October 1990		20 DEC 1990
International Searching Authority <sup>1</sup>		Signature of Authorized Officer <sup>20</sup>
ISA/US		 SCOTT A. CHAMBERS

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, <sup>16</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No <sup>18</sup>
X Y	WO, A, 89/01050 (BURG ET AL.) 09 February 1989, see entire document.	1,5,7,23,33,35, 40,42,48,71,83, 86 2-4,6,8-22,24- 32,34,36-39,41, 43-47,49-70,72- 82,84,85,87-92
X Y	WO, A, 88/10315 (Gingeras et al) 29 December 1988, see entire document.	1,5,7,11,12,19, 23,33,35,40,42, 48,49,71,83,86 2-4,6,8-10,13- 18,20-22,24-32, 34,36-39,41,43- 47,50-70,72-82, 84,85,87-92